

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
3 March 2005 (03.03.2005)

PCT

(10) International Publication Number  
**WO 2005/018637 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/407**,  
C07D 495/04, A61P 3/12

(21) International Application Number:  
PCT/GB2004/003546

(22) International Filing Date: 18 August 2004 (18.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0319759.7 22 August 2003 (22.08.2003) GB

(71) Applicant (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CY, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, SZ, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW only): **ASTRAZENECA AB** [SE/SE]; Sodertalje, S-SE-151 85 (SE).

(71) Applicant (for MG only): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London Greater London W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BENNETT, Stuart, Norman, Lile** [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield Cheshire SK10 4TG (GB). **SIMPSON, Iain** [GB/GB]; AstraZeneca R & D Alderley,

Alderley Park, Macclesfield Cheshire SK10 4TG (GB). **WHITTAMORE, Paul, Robert, Owen** [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield Cheshire SK10 4TG (GB).

(74) Agent: **ASTRAZENECA**; Global Intellectual Property, S-SE-151 85 Sodertalje (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLIC AMIDE DERIVATIVES WHICH POSSESS GLYCOGEN PHOSPHORYLASE INHIBITORY ACTIVITY



(57) Abstract: A compound of the formula (1) or a pharmaceutically-acceptable salt, or pro-drug thereof; (1) wherein, for example: R<sup>4</sup> and R<sup>5</sup> together are either -S-C(R<sup>6</sup>=C(R<sup>7</sup>)- or -C(R<sup>7</sup>)=C(R<sup>6</sup>)-S-; R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen and halo; A is phenylene or heteroarylene; n is 0, 1 or 2; R<sup>1</sup> is halo, cyano or carboxy; R<sup>2</sup> is for example methyl; R<sup>3</sup> is for example selected from halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (optionally substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl; possess glycogen phosphorylase inhibitory activity and accordingly have value in the treatment of disease states associated with increased glycogen phosphorylase activity. Processes for the manufacture of compounds and pharmaceutical compositions containing them are described.

WO 2005/018637 A1

HETEROCYCLIC AMIDE DERIVATIVES WHICH POSSESS GLYCOGEN  
PHOSPHORYLASE INHIBITORY ACTIVITY

The present invention relates to heterocyclic amide derivatives, pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof. These heterocyclic amides possess  
5 glycogen phosphorylase inhibitory activity and accordingly have value in the treatment of disease states associated with increased glycogen phosphorylase activity and thus are potentially useful in methods of treatment of a warm-blooded animal such as man. The invention also relates to processes for the manufacture of said heterocyclic amide derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of  
10 medicaments to inhibit glycogen phosphorylase activity in a warm-blooded animal such as man.

The liver is the major organ regulating glycaemia in the post-absorptive state. Additionally, although having a smaller role in the contribution to post-prandial blood glucose levels, the response of the liver to exogenous sources of plasma glucose is key to an ability to  
15 maintain euglycaemia. An increased hepatic glucose output (HGO) is considered to play an important role in maintaining the elevated fasting plasma glucose (FPG) levels seen in type 2 diabetics; particularly those with a FPG >140mg/dl (7.8mM). (Weyer et al, (1999), J Clin Invest 104: 787-794; Clore & Blackgard (1994), Diabetes 43: 256-262; De Fronzo, R. A., et al, (1992) Diabetes Care 15; 318 - 355; Reaven, G.M. (1995) Diabetologia 38; 3-13).

20 Since current oral, anti-diabetic therapies fail to bring FPG levels to within the normal, non-diabetic range and since raised FPG (and glycoHbA1c) levels are risk factors for both macro- (Charles, M.A. et al (1996) Lancet 348, 1657-1658; Coutinho, M. et al (1999) Diabetes Care 22; 233-240; Shaw, J.E. et al (2000) Diabetes Care 23, 34-39) and micro-vascular disease (DCCT Research Group (1993) New. Eng. J. Med. 329; 977-986); the  
25 reduction and normalisation of elevated FPG levels remains a treatment goal in type 2 DM.

It has been estimated that, after an overnight fast, 74% of HGO was derived from glycogenolysis with the remainder derived from gluconeogenic precursors (Hellerstein et al (1997) Am J Physiol, 272: E163). Glycogen phosphorylase is a key enzyme in the generation by glycogenolysis of glucose-1-phosphate, and hence glucose in liver and also in other tissues  
30 such as muscle and neuronal tissue.

Liver glycogen phosphorylase activity is elevated in diabetic animal models including the db/db mouse and the fa/fa rat (Aiston S et al (2000). Diabetologia 43, 589-597).

Inhibition of hepatic glycogen phosphorylase with chloroindole inhibitors (CP91149 and CP320626) has been shown to reduce both glucagon stimulated glycogenolysis and glucose output in hepatocytes (Hoover et al (1998) J Med Chem 41, 2934-8; Martin et al (1998) PNAS 95, 1776-81). Additionally, plasma glucose concentration is reduced, in a dose  
5 related manner, db/db and ob/ob mice following treatment with these compounds.

Studies in conscious dogs with glucagon challenge in the absence and presence of another glycogen phosphorylase inhibitor, Bay K 3401, also show the potential utility of such agents where there is elevated circulating levels of glucagon, as in both Type 1 and Type 2 diabetes. In the presence of Bay R 3401, hepatic glucose output and arterial plasma glucose  
10 following a glucagon challenge were reduced significantly (Shiota et al, (1997), Am J Physiol, 273: E868).


The heterocyclic amides of the present invention possess glycogen phosphorylase inhibitory activity and accordingly are expected to be of use in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac  
15 ischaemia and obesity, particularly type 2 diabetes.

Our patent application WO 02/20530 discloses a spectrum of active glycogen phosphorylase inhibitors, amongst which are a very limited number of amino-indan containing compounds.

Our co-pending patent applications PCT/GB03/00883 and PCT/GB03/00875 disclose  
20 a variety of substituted amino-indan glycogen phosphorylase inhibitors, generally containing only one substituent on the nitrogen of the amino-indan moiety, although a number are disubstituted and contain an N-acetyl group as one substituent.

Surprisingly, we have found that a group of N-disubstituted amino-indans have improved physical properties (for example solubility, plasma-protein binding) in comparison  
25 with that of the compounds previously disclosed, which are particularly beneficial for a pharmaceutical.

According to one aspect of the present invention there is provided a compound of formula (1):



wherein:

$R^4$  and  $R^5$  together are either  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

- 5  $R^6$  and  $R^7$  are independently selected from hydrogen, halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy and (1-4C)alkanoyl;  
A is phenylene or heteroarylene;  
n is 0, 1 or 2;
- 10  $R^1$  is independently selected from halo, nitro, cyano, hydroxy, carboxy, carbamoyl, *N*-(1-4C)alkylcarbamoyl, *N,N*-((1-4C)alkyl)<sub>2</sub>carbamoyl, sulphamoyl, *N*-(1-4C)alkylsulphamoyl, *N,N*-((1-4C)alkyl)<sub>2</sub>sulphamoyl,  $-S(O)_b(1-4C)alkyl$  (wherein b is 0, 1, or 2),  $-OS(O)_2(1-4C)alkyl$ , (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkanoyloxy, hydroxy(1-4C)alkyl, fluoromethyl, difluoromethyl,
- 15 trifluoromethyl, trifluoromethoxy and  $-NH SO_2(1-4C)alkyl$ ;  
or, when n is 2, the two  $R^1$  groups, together with the carbon atoms of A to which they are attached, may form a 4 to 7 membered saturated ring, optionally containing 1 or 2 heteroatoms independently selected from O, S and N, and optionally being substituted by one or two methyl groups;
- 20 one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ;  
 $R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl (optionally substituted on alkyl with hydroxy), methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives
- 25 thereof, (amino)(hydroxy)(2-3C)alkyl, (aminocarbonyl)(hydroxy)(2-3C)alkyl, (methylaminocarbonyl)(hydroxy)(2-3C)alkyl, (dimethylaminocarbonyl)(hydroxy)(2-3C)alkyl, (methylcarbonylamino)(hydroxy)(2-3C)alkyl, (methylS(O)<sub>p</sub>)(hydroxy)(2-3C)alkyl (wherein p is 0, 1 or 2);

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (optionally substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);

- 10 wherein any alkyl or alkoxy group within any group in  $R_{Na}$  and  $R_{Nb}$  may also optionally be substituted on an available carbon atom with a hydroxy group (provided that said carbon atom is not already substituted by a group linked by a heteroatom);  
provided that if  $R^2$  is (1-3C)alkyl or (1-4C)alkyl then  $R^3$  is not (1-4C)alkyl or (1-3C)alkyl; or a pharmaceutically acceptable salt or pro-drug thereof.

- 15 In another aspect of the invention, there is provided a compound of the formula (1) or a pharmaceutically acceptable salt or pro-drug thereof wherein one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ; and  $R_{Na}$  and  $R_{Nb}$  are selected from:

$R_{Na}$ : (1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl,


- 20 methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;


- 25

and wherein  $R^1$ ,  $R^4$ , A, m and n are as hereinbefore defined.

- 30 It is to be understood that when A is heteroarylene, the bridgehead atoms joining ring A to the ring may be heteroatoms. Therefore, for example, the definition of



when A is heteroarylene encompasses the structures:



It is to be understood that where substituents contain two substituents on an alkyl chain, in which both are linked by a heteroatom (for example two alkoxy substituents, or an amino and a hydroxy substituent), then these two substituents are not substituents on the same carbon atom of the alkyl chain.

In another aspect, the invention relates to compounds of formula (1) as hereinabove defined or to a pharmaceutically acceptable salt.

In another aspect, the invention relates to compounds of formula (1) as hereinabove defined or to a pro-drug thereof. Suitable examples of pro-drugs of compounds of formula (1) are in-vivo hydrolysable esters of compounds of formula (1). Therefore in another aspect, the invention relates to compounds of formula (1) as hereinabove defined or to an in-vivo hydrolysable ester thereof.

It is to be understood that, insofar as certain of the compounds of formula (1) defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses glycogen phosphorylase inhibition activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to hereinafter.

Within the present invention it is to be understood that a compound of the formula (1) or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be

understood that the invention encompasses any tautomeric form which has glycogen phosphorylase inhibition activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings. The formulae drawings within this specification can represent only one of the possible tautomeric forms and it is to be understood that the  
5 specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been possible to show graphically herein.

It is also to be understood that certain compounds of the formula (1) and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which have glycogen  
10 phosphorylase inhibition activity.

It is also to be understood that certain compounds of the formula (1) may exhibit polymorphism, and that the invention encompasses all such forms which possess glycogen phosphorylase inhibition activity.

The present invention relates to the compounds of formula (1) as hereinbefore defined  
15 as well as to the salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of the compounds of formula (1) and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of the compounds of formula (1) as hereinbefore defined which are sufficiently basic to form such  
20 salts. Such acid addition salts include for example salts with inorganic or organic acids affording pharmaceutically acceptable anions such as with hydrogen halides (especially hydrochloric or hydrobromic acid of which hydrochloric acid is particularly preferred) or with sulphuric or phosphoric acid, or with trifluoroacetic, citric or maleic acid. Suitable salts include hydrochlorides, hydrobromides, phosphates, sulphates, hydrogen sulphates,  
25 alkylsulphonates, arylsulphonates, acetates, benzoates, citrates, maleates, fumarates, succinates, lactates and tartrates. In addition where the compounds of formula (1) are sufficiently acidic, pharmaceutically acceptable salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases include for example an alkali metal salt, such as a sodium or potassium salt, an  
30 alkaline earth metal salt such as a calcium or magnesium salt, an ammonium salt or for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the invention may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the invention. A prodrug may be used to alter or improve the physical and/or pharmacokinetic profile of the parent compound and can be formed when the parent compound contains a suitable group or  
5 substituent which can be derivatised to form a prodrug. Examples of pro-drugs include in-vivo hydrolysable esters of a compound of the invention or a pharmaceutically-acceptable salt thereof.

Various forms of prodrugs are known in the art, for examples see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in  
10 Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- 15 d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- e) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

An in-vivo hydrolysable ester of a compound of formula (1) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is cleaved in the  
20 human or animal body to produce the parent acid or alcohol.

Suitable pharmaceutically acceptable esters for carboxy include alkyl, (1-6C)alkoxymethyl esters for example methoxymethyl, (1-6C)alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, (3-8C)cycloalkoxycarbonyloxy(1-6C)alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for  
25 example 5-methyl-1,3-dioxolen-2-onylmethyl; and (1-6C)alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

Suitable pharmaceutically-acceptable esters for hydroxy include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and  $\alpha$ -acyloxyalkyl ethers and  
30 related compounds which as a result of the *in-vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of  $\alpha$ -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in-vivo* hydrolysable ester forming groups for hydroxy include (1-10C)alkanoyl, for example acetyl; benzoyl; phenylacetyl; substituted



benzoyl and phenylacetyl, (1-10C)alkoxycarbonyl (to give alkyl carbonate esters), for example ethoxycarbonyl; di-((1-4C))alkylcarbamoyl and *N*-(di-((1-4C))alkylaminoethyl)-*N*-((1-4C))alkylcarbamoyl (to give carbamates); di-((1-4C))alkylaminoacetyl and carboxyacetyl. Examples of ring substituents on phenylacetyl and benzoyl include aminomethyl, ((1-4C))alkylaminomethyl and di-(((1-4C))alkyl)aminomethyl, and morpholino or piperazino linked from a ring nitrogen atom via a methylene linking group to the 3- or 4- position of the benzoyl ring. Other interesting in-vivo hydrolysable esters include, for example,  $R^A C(O)O((1-6C))alkyl-CO-$ , wherein  $R^A$  is for example, benzyloxy-((1-4C))alkyl, or phenyl). Suitable substituents on a phenyl group in such esters include, for example, 4-((1-4C))piperazino-((1-4C))alkyl, piperazino-((1-4C))alkyl and morpholino(1-4C)alkyl.

In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups. However references to individual alkyl groups such as "propyl" are specific for the straight chain version only and references to individual branched-chain alkyl groups such as *t*-butyl are specific for the branched chain version only. For example, "(1-3C)alkyl" includes methyl, ethyl, propyl and isopropyl, "(1-4C)alkyl" includes methyl, ethyl, propyl, isopropyl and *t*-butyl and examples of "(1-6C)alkyl" include the examples of "(1-4C)alkyl" and additionally pentyl, 2,3-dimethylpropyl, 3-methylbutyl and hexyl. An analogous convention applies to other generic terms, for example "(2-4C)alkenyl" includes vinyl, allyl and 1-propenyl and examples of "(2-6C)alkenyl" include the examples of "(2-4C)alkenyl" and additionally 1-butenyl, 2-butenyl, 3-butenyl, 2-methylbut-2-enyl, 3-methylbut-1-enyl, 1-pentenyl, 3-pentenyl and 4-hexenyl. Examples of "(2-4C)alkynyl" includes ethynyl, 1-propynyl and 2-propynyl and examples of "C<sub>2-6</sub>alkynyl" include the examples of "(2-4C)alkynyl" and additionally 3-butyne, 2-pentyne and 1-methylpent-2-ynyl.

The term "hydroxy(1-3C)alkyl" includes hydroxymethyl, hydroxyethyl, hydroxypropyl and hydroxyisopropyl. The term "hydroxy(2-3C)alkyl" includes hydroxyethyl, hydroxypropyl and hydroxyisopropyl. The term "hydroxy(1-4C)alkyl" includes hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxyisopropyl and hydroxybutyl. The term "hydroxy(1-4C)alkyl" also includes hydroxycyclopropyl and hydroxycyclobutyl. The term "hydroxyethyl" includes 1-hydroxyethyl and 2-hydroxyethyl. The term "hydroxypropyl" includes 1-hydroxypropyl, 2-hydroxypropyl and 3-hydroxypropyl and an analogous convention applies to terms such as hydroxybutyl. The term "dihydroxy(2-3C)alkyl" includes dihydroxyethyl, dihydroxypropyl and dihydroxyisopropyl. The term "dihydroxy(2-4C)alkyl"

includes dihydroxyethyl, dihydroxypropyl, dihydroxyisopropyl and dihydroxybutyl. The term “dihydroxypropyl” includes 1,2-dihydroxypropyl, 2,3-dihydroxypropyl and 1,3-dihydroxypropyl. An analogous convention applies to terms such as dihydroxyisopropyl and dihydroxybutyl. The term dihydroxy(2-4C)alkyl is not intended to include structures which  
 5 are geminally disubstituted and thereby unstable.

The term “trihydroxy(3-4C)alkyl” includes 1,2,3-trihydroxypropyl and 1,2,3-trihydroxybutyl. . The term trihydroxy(3-4C)alkyl is not intended to include structures which are geminally di- or tri-substituted and thereby unstable.

The term “halo” refers to fluoro, chloro, bromo and iodo. The term “halo(1-3C)alkyl”  
 10 includes fluoromethyl, chloromethyl, fluoroethyl, fluoropropyl and chloropropyl. The term “halo(1-4C)alkyl” includes “halo(1-3C)alkyl” and additionally fluorobutyl. The term “dihalo(1-4C)alkyl” includes difluoromethyl and dichloromethyl. The term “dihalo(1-3C)alkyl” includes difluoromethyl and dichloromethyl. The term “trihalo(1-4C)alkyl” includes trifluoromethyl.

15 Examples of “5- and 6-membered cyclic acetals and mono- and di-methyl derivatives thereof” are:

1,3-dioxolan-4-yl, 2-methyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxolan-4-yl; 2,2-dimethyl-1,3-dioxan-4-yl; 2,2-dimethyl-1,3-dioxan-5-yl; 1,3-dioxan-2-yl.

Examples of “(1-4C)alkoxy” include methoxy, ethoxy, propoxy and isopropoxy.

20 Examples of “(1-6C)alkoxy” include the examples of “(1-4C)alkoxy” and additionally butyloxy, *t*-butyloxy, pentoxy and 1,2-(methyl)<sub>2</sub>propoxy. Examples of “(1-4C)alkanoyl” include formyl, acetyl and propionyl. Examples of “(1-6C)alkanoyl” include the example of “(1-4C)alkanoyl” and additionally butanoyl, pentanoyl, hexanoyl and 1,2-(methyl)<sub>2</sub>propionyl. Examples of “(1-4C)alkanoyloxy” are formyloxy, acetoxo and propionoxy. Examples of “(1-  
 25 6C)alkanoyloxy” include the examples of “(1-4C)alkanoyloxy” and additionally butanoyloxy, pentanoyloxy, hexanoyloxy and 1,2-(methyl)<sub>2</sub>propionyl. Examples of “*N*-((1-4C)alkyl)amino” include methylamino and ethylamino. Examples of “*N*-((1-6C)alkyl)amino” include the examples of “*N*-((1-4C)alkyl)amino” and additionally pentylamino, hexylamino and 3-methylbutylamino. Examples of “*N,N*-((1-4C)alkyl)<sub>2</sub>amino” include *N,N*-  
 30 (methyl)<sub>2</sub>amino, *N,N*-(ethyl)<sub>2</sub>amino and *N*-ethyl-*N*-methylamino. Examples of “*N,N*-((1-6C)alkyl)<sub>2</sub>amino” include the example of “*N,N*-((1-4C)alkyl)<sub>2</sub>amino” and additionally *N*-methyl-*N*-pentylamino and *N,N*-(pentyl)<sub>2</sub>amino. Examples of “*N*-((1-4C)alkyl)carbamoyl” are methylcarbamoyl and ethylcarbamoyl. Examples of “*N*-((1-6C)alkyl)carbamoyl” are the

examples of "*N*-((1-4C)alkyl)carbamoyl" and additionally pentylcarbamoyl, hexylcarbamoyl and 1,2-(methyl)<sub>2</sub>propylcarbamoyl. Examples of "*N,N*-((1-4C)alkyl)<sub>2</sub>carbamoyl" are *N,N*-(methyl)<sub>2</sub>carbamoyl, *N,N*-(ethyl)<sub>2</sub>carbamoyl and *N*-methyl-*N*-ethylcarbamoyl. Examples of "*N,N*-((1-6C)alkyl)<sub>2</sub>carbamoyl" are the examples of "*N,N*-((1-4C)alkyl)<sub>2</sub>carbamoyl" and additionally *N,N*-(pentyl)<sub>2</sub>carbamoyl, *N*-methyl-*N*-pentylcarbamoyl and *N*-ethyl-*N*-hexylcarbamoyl. Examples of "*N*-((1-4C)alkyl)sulphamoyl" are *N*-(methyl)sulphamoyl and *N*-(ethyl)sulphamoyl. Examples of "*N*-((1-6C)alkyl)sulphamoyl" are the examples of "*N*-((1-4C)alkyl)sulphamoyl" and additionally *N*-pentylsulphamoyl, *N*-hexylsulphamoyl and 1,2-(methyl)<sub>2</sub>propylsulphamoyl. Examples of "*N,N*-((1-4C)alkyl)<sub>2</sub>sulphamoyl" are

10 *N,N*-(methyl)<sub>2</sub>sulphamoyl, *N,N*-(ethyl)<sub>2</sub>sulphamoyl and *N*-(methyl)-*N*-(ethyl)sulphamoyl. Examples of "*N,N*-((1-6C)alkyl)<sub>2</sub>sulphamoyl" are the examples of "*N,N*-((1-4C)alkyl)<sub>2</sub>sulphamoyl" and additionally *N,N*-(pentyl)<sub>2</sub>sulphamoyl, *N*-methyl-*N*-pentylsulphamoyl and *N*-ethyl-*N*-hexylsulphamoyl.

Examples of "cyano(1-3C)alkyl" and "cyano(1-4C)alkyl" are cyanomethyl, cyanoethyl and

15 cyanopropyl. Examples of "(3-6C)cycloalkyl" include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of "(3-6C)cycloalkyl(1-4C)alkyl" include cyclopropylmethyl, cyclopropylethyl, cyclopropylpropyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl. Examples of "cyano(1-3C)alkyl" and "cyano(1-4C)alkyl" substituted with hydroxy include 1-(hydroxy)-2-(cyano)ethyl.

20 The term "amino(1-4C)alkyl" includes aminomethyl, aminoethyl, aminopropyl, aminoisopropyl and aminobutyl. The term "aminoethyl" includes 1-aminoethyl and 2-aminoethyl. The term "aminopropyl" includes 1-aminopropyl, 2-aminopropyl and 3-aminopropyl and an analogous convention applies to terms such as aminoethyl and aminobutyl.

25 Examples of "(1-4C)alkoxy(1-4C)alkoxy" are methoxymethoxy, ethoxymethoxy, ethoxyethoxy and methoxyethoxy. Examples of "hydroxy(1-4C)alkoxy" are hydroxyethoxy and hydroxypropoxy. Examples of "hydroxypropoxy" are 2-hydroxypropoxy and 3-hydroxypropoxy. Examples of "(1-4C)alkoxy(1-4C)alkyl" include methoxymethyl, ethoxymethyl, methoxyethyl, ethoxypropyl and propoxymethyl. Examples of "(1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl" include methoxymethoxymethyl, ethoxyethoxyethyl, ethoxymethoxymethyl, methoxyethoxymethyl, methoxymethoxyethyl, methoxyethoxyethyl and ethoxymethoxymethyl. Examples of "di[(1-4C)alkoxy](2-4C)alkyl" include 1,2-dimethoxyethyl, 2,3-dimethoxypropyl and 1-methoxy-2-ethoxy-ethyl. Examples of

30

“(hydroxy)[(1-4C)alkoxy](2-4C)alkyl” include 1-hydroxy-2-methoxyethyl and 1-hydroxy-3-methoxypropyl.

Examples of “-S(O)<sub>b</sub>(1-4C)alkyl (wherein b is 0, 1 or 2)” include methylthio, ethylthio, propylthio, methylsulphinyl, ethylsulphinyl, propanesulphinyl, mesyl, ethylsulphonyl, 5 propylsulphonyl and isopropylsulphonyl.

Examples of “(1-6C)alkoxycarbonyl” include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl.

Examples of “(amino)(hydroxy)(2-3C)alkyl” and “(amino)(hydroxy)(2-4C)alkyl” include 1-amino-2-hydroxyethyl, 1-hydroxy-2-aminoethyl, 1-hydroxy-2-aminopropyl and 1-10 amino-2-hydroxypropyl. Examples of “(aminocarbonyl)(hydroxy)(2-3C)alkyl” and “(aminocarbonyl)(hydroxy)(2-4C)alkyl” include 1-(hydroxy)-2-(aminocarbonyl)ethyl and 1-(hydroxy)-3-(aminocarbonyl)propyl. Examples of “((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl” and “(methylaminocarbonyl)(hydroxy)(2-3C)alkyl” include 1-(hydroxy)-2-(*N*-methylaminocarbonyl)ethyl. Examples of “(di(1-4C)alkylaminocarbonyl)(hydroxy)(2-15 4C)alkyl” and “(dimethylaminocarbonyl)(hydroxy)(2-3C)alkyl” include 1-(hydroxy)-2-(*N,N*-dimethylaminocarbonyl)ethyl. Examples of “(1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl” and “methylcarbonylamino)(hydroxy)(2-3C)alkyl” include 1-hydroxy-2-(methylcarbonylamino)ethyl and 1-(methylcarbonylamino)-2-(hydroxy)ethyl.

Examples of “((1-4C)alkylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl” and “(methylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl” (wherein p is 0, 1 or 2) include 1-(hydroxy)-2-(methylthio)ethyl, 20 1-(hydroxy)-2-(methylsulfinyl)ethyl and 1-(hydroxy)-2-(methylsulfonyl)ethyl.

Examples of additional substitution on an alkyl or alkoxy group within a definition of R<sub>NA</sub> and R<sub>NB</sub> by hydroxy is to be understood to mean, for example, substitution of a hydroxy in di(halo)(1-4C)alkyl to give groups such as 1-hydroxy-2,2-difluoromethyl; or for example 25 substitution of a hydroxy into an (amino)(hydroxy)(2-4C)alkyl group to give a group such as 1,2-dihydroxy-3-aminopropyl; or for example substitution of a hydroxy into a “((1-4C)alkylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl, to give for example HOCH<sub>2</sub>CH<sub>2</sub>S(O)<sub>2</sub>CH<sub>2</sub>CH(OH)-, or C<sub>2</sub>H<sub>5</sub>S(O)<sub>2</sub>CH<sub>2</sub>CH(OH)CH(OH)-.

Within this specification composite terms are used to describe groups comprising 30 more than one functionality such as -(1-4C)alkylSO<sub>2</sub>(1-4C)alkyl. Such terms are to be interpreted in accordance with the meaning which is understood by a person skilled in the art for each component part. For example -(1-4C)alkylSO<sub>2</sub>(1-4C)alkyl includes

-methylsulphonylmethyl, -methylsulphonylethyl, -ethylsulphonylmethyl, and -propylsulphonylbutyl.

Where optional substituents are chosen from "0, 1, 2 or 3" groups it is to be understood that this definition includes all substituents being chosen from one of the specified  
 5 groups or the substituents being chosen from two or more of the specified groups. An analogous convention applies to substituents chose from "0, 1 or 2" groups and "1 or 2" groups.

"Heteroarylene" is a diradical of a heteroaryl group. A heteroaryl group is an aryl, monocyclic ring containing 5 to 7 atoms of which 1, 2, 3 or 4 ring atoms are chosen from  
 10 nitrogen, sulphur or oxygen. Examples of heteroarylene are oxazolylene, oxadiazolylene, pyridylene, pyrimidinylene, imidazolylene, triazolylene, tetrazolylene, pyrazinylene, pyridazinylene, pyrrolylene, thienylene and furylene.

Suitable optional substituents for heteroaryl groups, unless otherwise defined, are 1, 2 or 3 substituents independently selected from halo, cyano, nitro, amino, hydroxy, (1-4C)alkyl,  
 15 (1-4C)alkoxy, (1-4C)alkylS(O)<sub>b</sub> (wherein b is 0, 1 or 2), *N*-((1-4C)alkyl)amino and *N,N*-((1-4C)alkyl)<sub>2</sub>amino. Further suitable optional substituents for "heteroaryl" groups are 1, 2 or 3 substituents independently selected from fluoro, chloro, cyano, nitro, amino, methylamino, dimethylamino, hydroxy, methyl, ethyl, methoxy, methylthio, methylsulfinyl and methylsulfonyl.

20 Preferred values of A, R<sup>1</sup> to R<sup>7</sup> and n are as follows. Such values may be used where appropriate with any of the definitions, claims, aspects or embodiments defined hereinbefore or hereinafter.

In one embodiment of the invention are provided compounds of formula (1), in an alternative embodiment are provided pharmaceutically-acceptable salts of compounds of  
 25 formula (1), in a further alternative embodiment are provided in-vivo hydrolysable esters of compounds of formula (1), and in a further alternative embodiment are provided pharmaceutically-acceptable salts of in-vivo hydrolysable esters of compounds of formula (1).

In one aspect of the present invention there is provided a compound of formula (1) as depicted above wherein R<sup>4</sup> and R<sup>5</sup> are together -S-C(R<sup>6</sup>)=C(R<sup>7</sup>)-.

30 In another aspect of the invention R<sup>4</sup> and R<sup>5</sup> are together -C(R<sup>7</sup>)=C(R<sup>6</sup>)-S-.

In a further aspect of the invention, R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, halo or (1-6C)alkyl.

Preferably  $R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl.

Particularly  $R^6$  and  $R^7$  are independently selected from hydrogen or chloro.

More particularly one of  $R^6$  and  $R^7$  is chloro.

5 In one embodiment, one of  $R^6$  and  $R^7$  is chloro and the other is hydrogen.

In another embodiment, both  $R^6$  and  $R^7$  are chloro.

In one aspect of the invention A is phenylene.

In another aspect of the invention A is heteroarylene.


10 Preferably A is selected from phenylene, pyridylene, pyrimidinylene, pyrrolylene, imidazolylen, triazolylen, tetrazolylen, oxazolylen, oxadiazolylen, thienylene and furylene.

Further suitable values for A are phenylene, pyridylene, pyrimidinylene, pyrrolylene and imidazolylen.

Further suitable values for A are phenylene, pyridylene and pyrimidinylene.

15 Further suitable values for A are phenylene and pyridylene.

In one embodiment, when A is heteroarylene, there is a nitrogen in a bridgehead position. In another embodiment, when A is heteroarylene, the heteroatoms are not in bridgehead positions. It will be appreciated that the preferred (more stable) bridgehead position is as shown below:



20

In one aspect of the invention n is 0 or 1.

In one aspect preferably n is 1.

In another aspect, preferably n is 0.

25 When n is 2, and the two  $R^1$  groups, together with the carbon atoms of A to which they are attached, form a 4 to 7 membered saturated ring, optionally containing 1 or 2 heteroatoms independently selected from O, S and N, conveniently such a ring is a 5 or 6 membered ring. In one embodiment, such a 5 or 6 membered ring contains two O atoms (ie a cyclic acetal). When the two  $R^1$  groups together form such a cyclic acetal, preferably it is not substituted. Most preferably the two  $R^1$  groups together are the group  $-O-CH_2-O-$ .

In another aspect of the present invention  $R^1$  is selected from halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl and (1-4C)alkoxy.

In a further aspect  $R^1$  is selected from halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl,  $-S(O)_b(1-4C)alkyl$  (wherein b is 0, 1 or 2),  $-OS(O)_2(1-4C)alkyl$ , (1-4C)alkyl and (1-4C)alkoxy.

In a further aspect  $R^1$  is selected from halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl,  $-S(O)_bMe$  (wherein b is 0, 1 or 2),  $-OS(O)_2Me$ , methyl and methoxy.

In a further aspect,  $R^1$  is (1-4C)alkyl.

10 Preferably  $R^1$  is selected from halo and (1-4C)alkoxy.

In another embodiment preferably  $R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ .

In one aspect  $R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from:

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl,  
15 dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl (optionally substituted on alkyl with hydroxy),  
methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl,  
(hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives  
thereof, (amino)(hydroxy)(2-3C)alkyl, (aminocarbonyl)(hydroxy)(2-3C)alkyl,  
(methylaminocarbonyl)(hydroxy)(2-3C)alkyl, (dimethylaminocarbonyl)(hydroxy)(2-3C)alkyl,  
20 (methylcarbonylamino)(hydroxy)(2-3C)alkyl, (methylS(O)<sub>p</sub>)(hydroxy)(2-3C)alkyl (wherein  
p is 0, 1 or 2);

and  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl,  
25 dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (optionally  
substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl,  
di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-  
membered acetals and mono- and di-methyl derivatives thereof, (amino)(hydroxy)(2-4C)alkyl,  
(aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl,  
(di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl,  
30 ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);

provided that when  $R_{Na}$  is (1-3C)alkyl, then  $R_{Nb}$  is not (1-4C)alkyl.

In another aspect  $R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(2-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

5 and  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-  
10 methyl derivatives thereof;  
provided that when  $R_{Na}$  is (1-3C)alkyl, then  $R_{Nb}$  is not (1-4C)alkyl.

In another aspect  $R^3$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl,  
15 methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

and  $R^2$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(2-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl,  
20 (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;  
provided that when  $R_{Na}$  is (1-3C)alkyl, then  $R_{Nb}$  is not (1-4C)alkyl.

In another aspect  $R^2$  is selected from  $R_{Na}$  and  $R^3$  is selected from  $R_{Nb}$ , wherein  $R_{Na}$   
25 and  $R_{Nb}$  are selected from any of the values for these groups defined hereinbefore or hereinafter.

In one embodiment, any alkyl or alkoxy group within any group in  $R_{Na}$  and  $R_{Nb}$  is additionally substituted on an available carbon atom with a hydroxy group (provided that said carbon atom is not already substituted by a group linked by a heteroatom).

30 In another embodiment, any alkyl or alkoxy group within any group in  $R_{Na}$  and  $R_{Nb}$  is not additionally substituted on an available carbon atom with a hydroxy group.

In one aspect,  $R_{Na}$  is selected from (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3C)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl and cyano(1-3C)alkyl.



In one embodiment  $R_{Na}$  is selected from methyl, ethyl, fluoromethyl, chloromethyl, dichloromethyl, difluoromethyl, trifluoromethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxy ethyl, dihydroxypropyl and cyanomethyl.

In another aspect  $R_{Na}$  is selected from (1-4C)alkyl, hydroxy(1-4C)alkyl, and  
5 (1-4C)alkoxy(1-4C)alkyl.

In another embodiment  $R_{Na}$  is selected from:  
(1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3C)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl and (hydroxy)(methoxy)ethyl.

10 In another embodiment  $R_{Na}$  is selected from:  
methyl, ethyl, fluoromethyl, difluoromethyl, trifluoromethyl, hydroxymethyl, hydroxyethyl, dihydroxyethyl, dihydroxypropyl, methoxymethyl, methoxyethyl and dimethoxyethyl.

In another embodiment  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl, hydroxyethyl, dihydroxyethyl, and dihydroxypropyl.

15 In another embodiment  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl.

In another embodiment  $R_{Na}$  is selected from methyl and hydroxyethyl.

In another embodiment  $R_{Na}$  is selected from methyl and ethyl.

In another embodiment  $R_{Na}$  is methyl.

20 In one embodiment  $R_{Nb}$  is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(1-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(1-4C)alkyl, and ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(1-4C)alkyl (wherein p is 0, 1 or 2).

In another embodiment  $R_{Nb}$  is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof.

In another embodiment  $R_{Nb}$  is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl,

(hydroxy)[(1-4C)alkoxy](1-4C)alkyl, (amino)(hydroxy)(1-4C)alkyl, (aminocarbonyl)(hydroxy)(1-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(1-4C)alkyl, and ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(1-4C)alkyl  
 5 (wherein p is 0, 1 or 2).

In another embodiment R<sub>Nb</sub> is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof.

In another embodiment R<sub>Nb</sub> is selected from hydroxy(1-4C)alkyl and dihydroxy(2-4C)alkyl.  
 10 4C)alkyl.

In another embodiment, R<sub>Nb</sub> is selected from dihydroxy(2-4C)alkyl and (hydroxy)[(1-4C)alkoxy](1-4C)alkyl.

In one aspect R<sub>Nb</sub> is selected from hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1,2,3-  
 15 trihydroxypropyl, methoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, hydroxyethoxyethyl, ,3-dioxolan-4-yl, 2-methyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxolan-4-yl; 2,2-dimethyl-1,3-dioxan-4-yl; 2,2-dimethyl-1,3-dioxan-5-yl; 1,3-dioxan-2-yl.

In another aspect R<sub>Nb</sub> is selected from hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl ,3-dioxolan-  
 20 4-yl, 2-methyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxan-4-yl, 2,2-dimethyl-1,3-dioxan-5-yl and 1,3-dioxan-2-yl.

In another aspect R<sub>Nb</sub> is selected from hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl and 1,3-dihydroxypropyl.

In a further aspect, R<sub>Nb</sub> is selected from hydroxymethyl, 1-hydroxyethyl, 2-  
 25 hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-(methylthio)ethyl, 1-(hydroxy)-2-(methylsulfonyl)ethyl, 1-(hydroxy)-2-(cyano)ethyl, 1-(hydroxy)-2-(amino)ethyl, 1-(amino)-2-(hydroxy)ethyl, 1-(hydroxy)-2-(aminocarbonyl)ethyl, 1-(hydroxy)-3-(aminocarbonyl)propyl, 1-(hydroxy)-2-(N-methylaminocarbonyl)ethyl, 1-  
 30 (hydroxy)-2-(N,N-dimethylaminocarbonyl)ethyl and 1-(methylcarbonylamino)-2-(hydroxy)ethyl.

In a further aspect, R<sub>Nb</sub> is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-

dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-(methylthio)ethyl, 1-(hydroxy)-2-(methylsulfonyl)ethyl, 1-(hydroxy)-2-(cyano)ethyl, 1-(hydroxy)-2-(amino)ethyl and 1-(amino)-2-(hydroxy)ethyl.

In a further aspect,  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-(methylthio)ethyl and 1-(hydroxy)-2-(methylsulfonyl)ethyl.

In a further aspect,  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl and 1-(hydroxy)-2-(methoxy)ethyl.

In a further aspect,  $R_{Nb}$  is selected from 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl and 1-(hydroxy)-2-(methoxy)ethyl.

In one aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

n is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

$R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(2-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

and  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

provided that when  $R_{Na}$  is (1-3C)alkyl, then  $R_{Nb}$  is not (1-4C)alkyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein A is heteroarylene;

n is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

$R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

5  $R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(2-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

10 and  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-

15 methyl derivatives thereof;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In one aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0, 1 or 2;

20  $R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

$R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

$R^3$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

and  $R^2$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from:

$R_{Nb}$ : (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl,

30 hydroxy(2-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

provided that when R<sub>Na</sub> is (1-3C)alkyl, then R<sub>Nb</sub> is not (1-4C)alkyl;  
and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein  
A is heteroaryl;

5 n is 0, 1 or 2;

R<sup>4</sup> and R<sup>5</sup> are together -S-C(R<sup>6</sup>)=C(R<sup>7</sup>)- or -C(R<sup>7</sup>)=C(R<sup>6</sup>)-S-;

R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, chloro, bromo or methyl;

R<sup>1</sup> is selected from fluoro, chloro, methyl, ethyl, methoxy and -O-CH<sub>2</sub>-O-;

R<sup>3</sup> is selected from R<sub>Na</sub> where R<sub>Na</sub> is selected from

10 R<sub>Na</sub>: (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3C)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, (hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

and R<sup>2</sup> is selected from R<sub>Nb</sub> where R<sub>Nb</sub> is selected from:

15 R<sub>Nb</sub>: (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(2-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

20 and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0, 1 or 2;

R<sup>4</sup> and R<sup>5</sup> are together -S-C(R<sup>6</sup>)=C(R<sup>7</sup>)- or -C(R<sup>7</sup>)=C(R<sup>6</sup>)-S-;

25 R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, chloro, bromo or methyl;

R<sup>1</sup> is selected from fluoro, chloro, methyl, ethyl, methoxy and -O-CH<sub>2</sub>-O-;

one of R<sup>2</sup> and R<sup>3</sup> is selected from R<sub>Na</sub>, and the other is selected from R<sub>Nb</sub>;

R<sub>Na</sub> is selected from: (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3C)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl, dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl, methoxymethyl,

30 ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl and (hydroxy)(methoxy)ethyl;

R<sub>Nb</sub> is selected from: hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](1-4C)alkyl,

(hydroxy)[(1-4C)alkoxy](1-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

5 A is phenylene;

n is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

$R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

10 one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ;

$R_{Na}$  is selected from: methyl, ethyl, fluoromethyl, difluoromethyl, trifluoromethyl, hydroxymethyl, hydroxyethyl, dihydroxyethyl, dihydroxypropyl, methoxymethyl, methoxyethyl and dimethoxyethyl.

$R_{Nb}$  is selected from: hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-

15 dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1,2,3-trihydroxypropyl, methoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl, hydroxyethoxyethyl, ,3-dioxolan-4-yl, 2-methyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxolan-4-yl; 2,2-dimethyl-1,3-dioxan-4-yl; 2,2-dimethyl-1,3-dioxan-5-yl; 1,3-dioxan-2-yl; and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

20 In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

n is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

25  $R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ;

$R_{Na}$  is selected from: methyl, ethyl, hydroxymethyl, hydroxyethyl, dihydroxyethyl, and dihydroxypropyl;

$R_{Nb}$  is selected from: hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-

30 dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, ,3-dioxolan-4-yl, 2-methyl-1,3-dioxolan-4-yl, 2,2-dimethyl-1,3-dioxolan-4-yl; 2,2-dimethyl-1,3-dioxan-4-yl; 2,2-dimethyl-1,3-dioxan-5-yl and 1,3-dioxan-2-yl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ; .

5  $R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is (1-4C)alkyl;

one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ;

$R_{Na}$  is selected from: methyl and ethyl;

$R_{Nb}$  is selected from hydroxymethyl, hydroxyethyl, hydroxypropyl, dihydroxyethyl, 1,2-

10 dihydroxypropyl, 2,3-dihydroxypropyl, and 1,3-dihydroxypropyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0, 1 or 2;

15  $R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, chloro, bromo or methyl;

$R^1$  is selected from fluoro, chloro, methyl, ethyl, methoxy and  $-O-CH_2-O-$ ;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

20  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (optionally substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives

25 thereof, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

30 In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, chloro, bromo or methyl;

R<sup>1</sup> is selected from fluoro, chloro, methyl, ethyl, methoxy and -O-CH<sub>2</sub>-O-;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

5 R<sup>3</sup> is selected from R<sub>Nb</sub> where R<sub>Nb</sub> is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, and ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2); and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

15 A is phenylene;

$n$  is 0, 1 or 2;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, chloro, bromo or methyl;

R<sup>1</sup> is selected from fluoro, chloro, methyl, ethyl, methoxy and -O-CH<sub>2</sub>-O-;

20 R<sup>2</sup> is selected from R<sub>Na</sub> where R<sub>Na</sub> is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

R<sup>3</sup> is selected from R<sub>Nb</sub> where R<sub>Nb</sub> is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, (amino)(hydroxy)(2-4C)alkyl,

25 (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, and ((1-4C)alkylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);

30

In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

n is 0;



$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

- 5  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, and ((1-4C)alkylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

- 15 n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

- 20  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, cyano(1-4C)alkyl (substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, (amino)(hydroxy)(2-4C)alkyl and ((1-4C)alkylS(O)<sub>p</sub>)-(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);
- and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

25

In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

- 30  $R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

$R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-(methylthio)ethyl, 1-(hydroxy)-2-(methylsulfonyl)ethyl, 1-(hydroxy)-2-(cyano)ethyl, 1-(hydroxy)-2-(amino)ethyl, 1-(amino)-2-(hydroxy)ethyl, 1-(hydroxy)-2-(aminocarbonyl)ethyl, 1-(hydroxy)-3-(aminocarbonyl)propyl, 1-(hydroxy)-2-(N-methylaminocarbonyl)ethyl, 1-(hydroxy)-2-(N,N-dimethylaminocarbonyl)ethyl, 1-(methylcarbonylamino)-2-(hydroxy)ethyl; and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

10 In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

15  $R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

$R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-

20 (methylthio)ethyl, 1-(hydroxy)-2-(methylsulfonyl)ethyl, 1-(hydroxy)-2-(cyano)ethyl, 1-(hydroxy)-2-(amino)ethyl and 1-(amino)-2-(hydroxy)ethyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

25 n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

30  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl, 1-(hydroxy)-2-(methoxy)ethyl, 1-(hydroxy)-2-(methylthio)ethyl and 1-(hydroxy)-2-(methylsulfonyl)ethyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein A is phenylene;

n is 0;

5  $R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

$R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from hydroxymethyl, 1-hydroxyethyl, 2-

10 hydroxyethyl, hydroxypropyl, hydroxyisobutyl, dihydroxyethyl, 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl and 1-(hydroxy)-2-(methoxy)ethyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

In another aspect of the invention is provided a compound of the formula (I) wherein

A is phenylene;

15 n is 0;

$R^4$  and  $R^5$  are together  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;


$R^6$  and  $R^7$  are independently selected from hydrogen or chloro;

$R^2$  is selected from  $R_{Na}$  where  $R_{Na}$  is selected from methyl, ethyl, hydroxymethyl and hydroxyethyl;

20  $R^3$  is selected from  $R_{Nb}$  where  $R_{Nb}$  is selected from 1,2-dihydroxypropyl, 2,3-dihydroxypropyl, 1,3-dihydroxypropyl and 1-(hydroxy)-2-(methoxy)ethyl;

and pharmaceutically acceptable salts and in-vivo hydrolysable esters thereof.

Preferred compounds of the invention are of the formula (1A), wherein  $R^1$  to  $R^5$  and n are as defined in any aspect or embodiment described hereinbefore or hereinafter.



(1A)


In one aspect, preferred compounds of the invention are compounds of the formula (1) or (1A) as defined hereinbefore or hereinafter wherein  $R^3$  contains an hydroxy group on the carbon adjacent to the carbonyl group. Further preferred compounds of the invention are

compounds of the formula (1) or (1A) as defined hereinbefore or hereinafter wherein  $R^3$  contains an amino group on the carbon adjacent to the carbonyl group.

Particular compounds of the invention are each of the Examples, each of which provides a further independent aspect of the invention. In a further aspect of the invention there is provided any two or more of the Examples or a pharmaceutically acceptable salt or pro-drug thereof.


Another aspect of the present invention provides a process for preparing a compound of formula (1) or a pharmaceutically acceptable salt or an in-vivo hydrolysable ester thereof which process (wherein A,  $R^1$  to  $R^5$  and n are, unless otherwise specified, as defined in formula (1)) comprises of:

a) reacting an acid of the formula (2):



(2)

or an activated derivative thereof; with an amine of formula (3):



(3)

and thereafter if necessary:

- i) converting a compound of the formula (1) into another compound of the formula (1);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or in-vivo hydrolysable ester.

Specific reaction conditions for the above reaction are as follows.


*Process a)* Acids of formula (2) and amines of formula (3) may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, or for example carbonyldiimidazole, 1-ethyl-3-(3-dimethylaminopropyl)carbodi-imide hydrochloride (EDCI) and dicyclohexyl-carbodiimide (DCCI), optionally in the presence of a catalyst such as 1-

hydroxybenzotriazole, dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for example triethylamine, di-isopropylethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran and

5 dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides, and active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the  
10 presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.


A compounds of formula (2) may be prepared according to Scheme 1:



Scheme 1

Compounds of formula (2a) are commercially available or they are known compounds or they are prepared by processes known in the art.

Compounds of formula (2b) may also be prepared as illustrated in Scheme 2:



Scheme 2


The conversion of compounds of formula (4) into compounds of formula (5) may be carried out by directed ortho lithiation reactions (J. Org. Chem, 2001, volume 66, 3662-3670), for example with n-butyl lithium and (CHO)N(alkyl)<sub>2</sub>. The protecting group P' in compounds of formula (4) must be suitable directing group for this reaction and may be for example –

25 CO<sub>2</sub>tBu. Reaction of compounds of formula (5) with LCH<sub>2</sub>CO<sub>2</sub>R where L is a leaving group,

and replacement of the protecting group P' with an alternative P'' (for example -COalkyl) according to standard processes, gives a compound of formula (6). This may be cyclised using a base, for example potassium carbonate or sodium methoxide.


Compounds of formula (3) may be prepared according to Scheme 3:

5




**Scheme 3**


Compounds of formula (3a) are commercially available or they are known compounds or they are prepared by processes known in the art. For example, starting from primary amines of formula (7), in which R is H or a suitable protecting group, one or both of  $R^2$  and/or  $R^3$  may be introduced by acylation, (for example reacting with acetoxyacetic acid and 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride -EDAC), alkylation, reductive alkylation, sulphonation or related processes, followed by O-deprotection when appropriate. Alternatively, one or both of  $R^2$  and/or  $R^3$  may be obtained by modification of functionality in groups previously thus introduced, by reduction, oxidation, hydrolysis (for example the conversion of an acetoxy group to a hydroxy group), nucleophilic displacement, amidation, or a related process, or a combination of these processes, followed by O-deprotection when appropriate. It will be appreciated that such modifications may include modifications which convert one compound of the formula (1) into another compound of the formula (1).



Amines of formula (3) may alternatively be obtained by applying the processes described for the preparation of compounds of formula (3a) to compounds of formula (8) in 5 which W is NH<sub>2</sub> or a nitrogen atom with one or two suitable protecting groups.




Alternatively, amines of formula (3) may also be prepared by the process in Scheme 3A. Compounds of formula A are commercially available or they are known compounds or they are prepared by processes known in the art. For example compound A can be converted 10 to the phthalamido-protected intermediate C under standard conditions (Step 1). Alkylation can then be performed under standard conditions (Step 2: NaH, MeI, DMA). Removal of the phthalamide then affords amine D (Step 3; hydrazine hydrate, EtOH).




Scheme 3A

Compounds of the formula (3) where  $r = 1$  and wherein A is heteroarylene can be prepared from suitably functionalised cycloalkyl fused heterocycles. For example, when A is 5 pyridine,



compounds of formula (3b) and (3c) may be prepared from the corresponding azaindanone regioisomer according to Scheme 4 :-






Scheme 4

Step 1 is performed on a compound known in the literature (*Jpn. Kokai Tokkyo Koho*, **1995**, 14. JP 07070136). Steps 2, 3, 4, 5, 6, 7 and 8 are performed using standard techniques

5 known in the art.


It will be appreciated that the bromoazaindanone isomers (21a, 21b and 21c) could



be converted to the corresponding heterocyclic version of (3) by the means described in

Scheme 4. The bromoazaindanone can be prepared from the corresponding azaindanone by

10 standard techniques known in the art. The azaindanone (22a, 22b, 22c) are known in the literature or they are prepared by processes known in the art.



The process described above and shown in Scheme 4 may also be applied to other six membered heterocycles containing more than one nitrogen.

It will be appreciated that, in a similar manner, compounds of the formula (3) wherein A is heteroarylene containing a bridgehead nitrogen can be prepared from the appropriate  
5 suitably functionalised cycloalkyl fused heterocycles.

It will be appreciated that the processes described above for formation and modification of  $-NR^2C(O)R^3$  may be applied similarly whether to make the compound of formula (3) before coupling to the acid of formula (2) or whether to the product of such a coupling.

10 It will be appreciated that certain of the various ring substituents in the compounds of the present invention, for example  $R^1$  may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions may convert one compound of the formula (1)  
15 into another compound of the formula (1). Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using  
20 concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogen group. Particular examples of modifications include the reduction of a nitro group to an amino group  
25 by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where  
30 protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley

and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.




A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

Certain intermediates in the preparation of a compound of the formula (1) are novel and form another aspect of the invention.

Compounds of the invention generally possess improved physical properties (for example solubility and/or plasma-protein binding) in comparison with those of the  
5 compounds previously disclosed, which are particularly beneficial for a pharmaceutical. In combination with glycogen phosphorylase inhibitory activity, such physical properties render the compounds of the invention particularly useful as pharmaceuticals.

For example, the compounds of the invention generally show improved solubility in comparison with the equivalent compounds where  $R^2$  is H. This effect is illustrated by the  
10 thermodynamic solubilities of Examples 8 and 17, and Reference Example 1 given in the table below.


Example No	Structure	Solubility ( $\mu$ M)	Activity ( $\mu$ M) IC <sub>50</sub>
8		176	0.07
27		50	0.11
Reference Eg 1		1.4	0.4 approx

For Reference Example 1, See WO 02/20530 example 154.

The thermodynamic solubility data for the compounds of the invention as given above may be measured by agitating the compound in 0.1 M phosphate at pH7.4 for 24hours, then  
15 analysis of the supernatant (for example by LCUV/MS) using a solution (for example in DMSO) of known concentration as the calibrant.

Plasma Protein binding may be measured using an equilibrium dialysis technique, whereby compound is added to 10% plasma giving a concentration of 20  $\mu$ M and dialysed with isotonic buffer for 18 hours at 37°C. The plasma and buffer solutions are analysed using LCUVMS and the first apparent binding constant for the compound derived. The binding  
 5 constant is then used to determine the % free in 100% plasma.

The binding constant derived from the dialysis experiment is based upon a model of 1:1 binding between compound and albumin.



$$K1 = \frac{[PD]}{[P] \times [D]}$$

where P = free protein, D = free drug, PD = drug protein complex, K1 = first apparent binding  
 10 constant.

As stated hereinbefore the compounds defined in the present invention possesses glycogen phosphorylase inhibitory activity. This property may be assessed by, for example, using the procedure set out below.

15

### Assay

The activity of the compounds is determined by measuring the inhibitory effect of the compounds on glycogen degradation, the production of glucose-1-phosphate from glycogen is monitored by the multienzyme coupled assay, as described in EP 0 846 464 A2, general  
 20 method of Pesce et al ( Pesce, M A, Bodourian, S H, Harris, R C, and Nicholson, J F (1977) Clinical Chemistry 23, 1171 - 1717). The reactions were in 384well microplate format in a volume of 50 $\mu$ l. The change in fluorescence due to the conversion of the co-factor NAD to NADH is measured at 340nm excitation, 465nm emission in a Tecan Ultra Multifunctional Microplate Reader. The reaction is in 50mM HEPES, 3.5mM KH<sub>2</sub>PO<sub>4</sub>, 2.5mM MgCl<sub>2</sub>,  
 25 2.5mM ethylene glycol-bis(b-aminoethyl ether) *N,N,N',N'*-tetraacetic acid, 100mM KCl, 8mM D-(+)-glucose pH7.2, containing 0.5mM dithiothreitol, the assay buffer solution. Human recombinant liver glycogen phosphorylase  $\alpha$  (hrl GPa) 20nM is pre-incubated in assay buffer solution with 6.25mM NAD, 1.25mg type III glycogen at 1.25 mg ml<sup>-1</sup> the reagent buffer, for 30 minutes. The coupling enzymes, phosphoglucomutase and glucose-6-phosphate

dehydrogenase (Sigma) are prepared in reagent buffer, final concentration 0.25 Units per well. 20  $\mu$ l of the hrl GPa solution is added to 10  $\mu$ l compound solution and the reaction started with the addition of 20  $\mu$ l coupling enzyme solution. Compounds to be tested are prepared in 10  $\mu$ l 5% DMSO in assay buffer solution, with final concentration of 1% DMSO in the assay.

- 5 The non-inhibited activity of GPa is measured in the presence of 10  $\mu$ l 5% DMSO in assay buffer solution and maximum inhibition measured in the presence of 5  $\text{mgs ml}^{-1}$  N-ethylmaleimide. After 6 hours at 30°C Relative Fluorescence Units (RFUs) are measured at 340nm excitation, 465nm emission .

The assay is performed at a test concentration of inhibitor of 10  $\mu$ M or 100  $\mu$ M.

- 10 Compounds demonstrating significant inhibition at one or both of these concentrations may be further evaluated using a range of test concentrations of inhibitor to determine an  $\text{IC}_{50}$ , a concentration predicted to inhibit the enzyme reaction by 50%.

Activity is calculated as follows:-

- % inhibition =  $(1 - (\text{compound RFUs} - \text{fully inhibited RFUs}) / (\text{non-inhibited rate RFUs} - \text{fully inhibited RFUs})) * 100$ .
- 15

Typical  $\text{IC}_{50}$  values for compounds of the invention when tested in the above assay are in the range 100  $\mu$ M to 1nM. For example, Example 1 was found to have an  $\text{IC}_{50}$  of 265nM and Example 8 to have an  $\text{IC}_{50}$  of 176nm.

The inhibitory activity of compounds was further tested in rat primary hepatocytes.

- 20 Rat hepatocytes were isolated by the collagenase perfusion technique, general method of Seglen (P.O. Seglen, Methods Cell Biology (1976) 13 29-83). Cells were cultured on Nunclon six well culture plates in DMEM (Dulbecco's Modified Eagle's Medium) with high level of glucose containing 10% foetal calf serum, NEAA (non essential amino acids), Glutamine, penicillin /streptomycin ((100units/100ug)/ml) for 4 to 6 hours. The hepatocytes were then
- 25 cultured in the DMEM solution without foetal calf serum and with 10nM insulin and 10nM dexamethasone. Experiments were initiated after 18-20 hours culture by washing the cells and adding Krebs-Henseleit bicarbonate buffer containing 2.5mM  $\text{CaCl}_2$  and 1% gelatin. The test compound was added and 5 minutes later the cells were challenged with 25nM glucagon. The Krebs-Henseleit solution was removed after 60 min incubation at 37°C , 95% $\text{O}_2$ /5% $\text{CO}_2$  and
- 30 the glucose concentration of the Krebs-Henseleit solution measured.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula (1), or a pharmaceutically

acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible  
5 powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing  
10 or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents. In one aspect, the compositions of the invention are a form suitable for  
15 oral dosage.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc;  
20 preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

25 Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form  
30 together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or

condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example  
5 heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-  
10 oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as  
15 beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or  
20 wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of  
25 oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and  
30 condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.



Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The compound of formula (1) will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route

of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

The inhibition of glycogen phosphorylase activity described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example, in order to prevent, delay or treat type 2 diabetes mellitus, the compounds of the present invention or their pharmaceutically acceptable salts may be administered in

combination with one or more of the following agent(s):

- 1) Insulin and insulin analogues;
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide), prandial glucose regulators (for example repaglinide, nateglinide) and glucokinase activators
- 3) Agents that improve incretin action (for example dipeptidyl peptidase IV inhibitors, GLP-1 agonists)
- 4) Insulin sensitising agents including PPAR $\gamma$  agonists (for example pioglitazone and rosiglitazone); and agents with combined PPAR $\alpha$  and gamma activity
- 5) Agents that modulate hepatic glucose balance (for example metformin, fructose 1, 6 bisphosphatase inhibitors, glycogen synthase kinase inhibitors, glucokinase activators)
- 6) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 7) Agents that prevent the reabsorption of glucose by the kidney (SGLT inhibitors)
- 8) Agents designed to treat the complications of prolonged hyperglycaemia (for example aldose reductase inhibitors)
- 9) Anti-obesity agents (for example sibutramine and orlistat);
- 10) Anti- dyslipidaemia agents such as, HMG-CoA reductase inhibitors (statins, eg pravastatin); PPAR $\alpha$  agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); bile acid absorption inhibitors (IBATi) and nicotinic acid and analogues (niacin and slow release formulations);

- 11) Antihypertensive agents such as,  $\beta$  blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); Calcium antagonists (eg. nifedipine); Angiotensin receptor antagonists (eg candesartan),  $\alpha$  antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 12) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and  
5 antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors);  
antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low  
molecular weight analogues, hirudin) and warfarin;
- 13) Agents which antagonise the actions of glucagon; and
- 14) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin)  
10 and steroidal anti-inflammatory agents (eg. cortisone).

According to a further aspect of the present invention there is provided a compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use in a method of treatment of a warm-blooded animal such as man  
15 by therapy.

According to an additional aspect of the invention there is provided a compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use as a medicament.

According to an additional aspect of the invention there is provided a compound of the  
20 formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore, for use as a medicament in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

According to this another aspect of the invention there is provided the use of a  
25 compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

According to this another aspect of the invention there is provided the use of a  
30 compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the treatment of type 2 diabetes in a warm-blooded animal such as man.

- 43 -

According to a further feature of this aspect of the invention there is provided a method of producing a glycogen phosphorylase inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1).

5        According to this further feature of this aspect of the invention there is provided a method of treating type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1).

10       According to this further feature of this aspect of the invention there is provided a method of treating type 2 diabetes in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1).

As stated above the size of the dose required for the therapeutic or prophylactic  
15 treatment of a particular cell-proliferation disease will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

In addition to their use in therapeutic medicine, the compounds of formula (1) and their pharmaceutically acceptable salts are also useful as pharmacological tools in the  
20 development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the  
25 invention described herein also apply.

### **Examples**


The invention will now be illustrated by the following examples in which, unless stated otherwise:

30 (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C and under an atmosphere of an inert gas such as argon;

- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mmHg) with a bath temperature of up to 60°C;
- (iii) chromatography means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates;
- (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- (vi) where given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio dimethyl sulphoxide (DMSO- $\delta_6$ ) as solvent unless otherwise indicated, other solvents (where indicated in the text) include deuterated chloroform  $\text{CDCl}_3$ ;
- (vii) chemical symbols have their usual meanings; SI units and symbols are used;
- (viii) reduced pressures are given as absolute pressures in Pascals (Pa); elevated pressures are given as gauge pressures in bars;
- (ix) solvent ratios are given in volume : volume (v/v) terms;
- (x) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionisation (CI) mode using a direct exposure probe; where indicated ionisation was effected by electron impact (EI), fast atom bombardment (FAB) or electrospray (ESP); values for m/z are given; generally, only ions which indicate the parent mass are reported and unless otherwise stated the value quoted is (M-H)<sup>+</sup>;
- (xi) The following abbreviations may be used:
- |    |       |   |
|----|-------|---|
| 25 | SM    | starting material;  |
|    | EtOAc | ethyl acetate;  |
|    | MeOH  | methanol;   |
|    | EtOH  | ethanol;  |
|    | DCM   | dichloromethane;  |
| 30 | HOBT  | 1-hydroxybenzotriazole;   |
|    | DIPEA | di-isopropylethylamine;   |
|    | EDCI  | 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide<br>hydrochloride; |

5	EDAC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride;
	Et <sub>2</sub> O/ether	diethyl ether;
	THF	tetrahydrofuran;
	DMF	<i>N, N</i> -dimethylformamide;
	HATU	<i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N, N, N', N'</i> -tetramethyluroniumhexafluorophosphate
10	EDAC	1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride
	TFA	Trifluoroacetic acid
	DMTMM	4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride
	DMA	<i>N, N</i> -dimethylacetamide

15 **Example 1: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(methoxyacetyl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



DIPEA (171  $\mu$ L, 1.0 mmol), HOBt (54 mg, 0.4 mmol), methoxyacetic acid (31  $\mu$ L, 0.4 mmol) and EDAC (96 mg, 0.5 mmol) were added to a suspension of 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**, 150 mg, 0.4 mmol) in anhydrous DCM (5 mL). The reaction was stirred at ambient temperature for approximately 6 h. The volatiles were removed by evaporation under reduced pressure, the residue dissolved in EtOAc (20 mL), washed with saturated aqueous NaHCO<sub>3</sub> (20 mL), water (2 x 10 mL) then brine (10 mL) and dried (MgSO<sub>4</sub>). The volatiles were removed by evaporation under reduced pressure and the residue triturated (EtOAc:hexane, 1:10), collected by filtration, washed with hexane (2 x 5 mL) and dried to give the title compound (126 mg, 75%) as a brown solid.

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO)  $\delta$ : 2.61 (s, 1.5H), 2.74 (s, 1.5H), 2.98 (m, 1H), 3.18 (dd, 1H), 3.19 (s, 1.5H), 3.22 (s, 1.5H), 3.99 (d, 0.5H), 4.13 (q, 1H), 4.31 (d, 0.5H), 4.79 (m, 1H), 5.41

(d, 0.5H), 6.09 (d, 0.5H), 7.12 (m, 6H), 8.52 (ap t, 1H), 11.81 (br s, 0.5H), 11.89 (br s, 0.5H); MS m/z 418, 420.

The following examples were prepared in a similar manner to **Example 1** using 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**) as the amine and the appropriate commercially available carboxylic acid as the coupling partner:

There is no Example with Example No 2.

**Example 3: 2-Chloro-*N*-{(1*R*,2*R*)-1-[[3-hydroxy-2-(hydroxymethyl)propanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**

**Example 4: Ethyl 3-[(1*R*,2*R*)-2-{[(2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl](methyl)amino]-3-oxopropanoate**

**Example 5: 2-[(1*R*,2*R*)-2-{[(2-Chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl](methyl)amino]-2-oxoethyl acetate**


**Example 6: 2-Chloro-*N*-{(1*R*,2*R*)-1-[glycoloyl(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**








**Example 7: 2-Chloro-*N*-{(1*R*,2*R*)-1-[glyceroyl(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**

**Example 8: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2,3-dihydroxypropanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**

**Example 9: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2,3-dihydroxypropanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**

25 Note: Example 4 is an example of a pro-drug of a compound containing a carboxylic acid group and Example 5 is an example of a pro-drug of a compound containing a hydroxy group.




Example	R	<sup>1</sup> H NMR (300 MHz, d <sub>6</sub> -DMSO)	M/z
3		δ: 2.61 (s, 1.5H), 2.87 (s, 1.5H), 2.99 (m, 1H), 3.18 (m, 2H), 3.41 (m, 1H), 3.61 (m, 2H), 4.44 (t, 0.5H), 4.49 (t, 0.5H), 4.66 (m, 1H), 4.80 (m, 1H), 5.74 (d, 0.5H), 6.18 (0.5H, d), 6.97 (m, 2H), 7.21 (m, 4H), 8.47 (m, 1H), 11.78 (br s, 0.5H), 11.84 (br s, 0.5H)	446, 448 (M-H) <sup>+</sup>
4		δ: 1.07 (t, 1.5H), 1.11 (t, 1.5H), 2.61 (s, 1.5H), 2.79 (s, 1.5H), 2.97 (m, 1H), 3.20 (dd, 1H), 3.42 (m, 1H), 3.71 (d, 0.5H), 3.68 (d, 0.5H), 4.00 (q, 1H), 4.06 (q, 1H), 4.79 (m, 1H), 5.41 (d, 0.5H), 6.11 (d, 0.5H), 7.11 (m, 6H), 8.53 (m, 1H), 11.82 (br s, 0.5H), 11.91 (br s, 0.5H)	460, 462
5		δ: 2.02(s, 1.5H), 2.08(s, 1.5H), 2.61(s, 1.5H), 2.77(s, 1.5H), 2.98(dd, 1H), 3.21(dd, 1H), 4.75(m, 2H), 4.97(m, 1H), 5.33(d, 0.5H), 6.05(d, 0.5H), 7.08(m, 3H), 7.25(m, 3H), 8.55(dd, 1H), 11.82(s, 0.5H), 11.9(s, 0.5H).	446, 448
6		δ: 2.65 (s, 1.5H), 2.74(s, 1.5H), 3.0(m, 1H), 3.21(m, 1H), 4.17(m, 2H), 4.45(t, 0.5H), 4.65(m, 0.5H), 4.8(m, 1H), 5.34(d, 0.5H), 6.14(d, 0.5H), 7.05(m, 2H), 7.17(d, 1H), 7.28(m, 3H), 8.55(m, 1H), 11.87(d, 1H).	402, 404
7		δ(CDCl <sub>3</sub> ): 2.92(m, 4H), 3.35(m, 2H), 3.95(m, 2H), 4.78(m, 3H), 6.24(dd, 1H), 6.6(m, 3H), 7.1(m, 3.5), 7.8(m, 0.5H), 10.42(s, 1H), 10.86(s, 0.5H).	456, 458
8		2.73(s, 1.5H), 2.97(s, 1.5H), 3.05(m, 1H), 3.3(m, 1H), 3.6(m, 2H), 4.75(m, 4H), 5.74(d, 0.5H), 6.18(d, 0.5H), 7.15(m, 3H), 7.35(m, 3H), 8.6(m, 1H), 11.83(s, 0.5H), 11.88(s, 0.5H)	456, 458
9		δ: 2.7(s, 1.5H), 2.95(s, 1.5H), 3.05(m, 1H), 3.31(m, 1H), 3.6(m, 2H), 4.5(m, 1.5H), 4.77(d, 0.5H), 4.9(m, 1.5H), 5.38(d, 0.5H), 5.82(d, 0.5H), 6.24(d, 0.5H), 7.13(m, 3H), 3.34(m, 3H), 8.62(d, 1H), 11.86(s, 1H)	456, 458



There is no Example with Example No 10.

**Example 11: 2-Chloro-*N*-{[(1*R*,2*R*)-1-[(3-hydroxypropanoyl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**




5

LiBH<sub>4</sub> (14 mg, 0.67 mmol) was added to a solution of ethyl 3-[[[(2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1*H*-inden-1-yl](methyl)amino]-3-oxopropanoate (**Example 4**; 103 mg, 0.22 mmol) at 0 °C. The reaction was warmed and stirred at ambient temperature for 3 hours. Saturated aqueous NH<sub>4</sub>Cl (10 mL) was added and the aqueous phase was extracted with EtOAc (3 x 20 mL). The combined organic phases were washed with HCl (2N, 20 mL), saturated aqueous NaHCO<sub>3</sub> (20 mL), water (20 mL) and brine (20 mL) then dried (MgSO<sub>4</sub>) and the solvent was removed under reduced pressure. The crude product was purified by flash column chromatography (SiO<sub>2</sub>, EtOAc eluent) to afford the title compound (62 mg, 67%) as a solid.

15 <sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 2.43 (m, 1H), 2.60 (s, 1.5H), 2.70 (m, 1H), 2.80 (s, 1.5H), 2.97 (dd, 1H), 3.19 (dd, 1H), 3.62 (m, 2H), 4.46 (t, 0.5H), 4.52 (t, 0.5H), 4.78 (m, 1H), 5.55 (d, 0.5H), 6.14 (d, 0.5H), 7.12 (m, 6H), 8.49 (d, 0.5H), 8.54 (d, 0.5H), 11.82 (br s, 0.5H), 11.89 (br s, 0.5H); MS m/z 418, 420.

20 **Example 12: 2-Chloro-*N*-{[(1*R*,2*R*)-1-[glycoloyl(2-hydroxyethyl)amino]-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



LiBH<sub>4</sub> (3.0ml, 2M in THF, 6.0 mmol) was added to a solution of ethyl *N*-[(acetyloxy)acetyl]-*N*-{[(1*R*,2*R*)-2-[[[(2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1*H*-


inden-1-yl)glycinate (**Intermediate 11**; 620 mg, 1.2 mmol) in dry THF (25 ml) at 5°C and stirred for 30mins. The reaction was allowed to warm to ambient temperature, stirred for 20 hours, HCl (1N, 10ml) added and the aqueous phase extracted with EtOAc (50ml). The combined organic phases were washed with water (20 mL) and brine (20 mL) then dried

5 (MgSO<sub>4</sub>) and the solvent was removed under reduced pressure. The crude product was purified by flash column chromatography (SiO<sub>2</sub>, EtOAc:hexane eluent) to afford the title compound (230 mg, 44%) as a foam.

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ: 2.9 (m, 2H), 3.37 (m, 3H), 4.25 (m, 3H), 4.75 (m, 3H), 5.35 (d, 0.6H), 5.59 (m, 0.4H), 7.0 (s, 1H), 7.2 (m, 5H), 8.54 (m, 1H), 11.84 (m, 1H); MS m/z

10 432, 434.

**Example 13: 2-Chloro-N-[(1R,2R)-1-[(2R)-2-hydroxypropanoyl](methylamino)-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-b]pyrrole-5-carboxamide**



15 To a solution of 2-chloro-N-[(1R,2R)-1-(methylamino)-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-b]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**; 191mg, 0.5 mmol), (R)-lactic acid (45 mg, 0.5 mmol) and DIPEA (171 μL, 1.0 mmol) in anhydrous DMA (5 mL) was added HOBT (68 mg, 0.5 mmol) and EDCI (120 mg, 0.625 mmol). The reaction was stirred at ambient temperature for approximately 24 h, water (25 mL) was added and the


20 resulting precipitate filtered. The solid was purified by flash column chromatography (EtOAc:hexane, 1:1) to give the title compound (110mg, 52%).

<sup>1</sup>H NMR (mixture of rotamers) δ: 1.1 (d, 1.5H), 1.22 (d, 1.5H), 2.64 (s, 1.5H), 2.83 (s, 1.5H), 3.0 (m, 1H), 3.2 (m, 1H), 4.5 (m, 1H), 4.8 (m, 2H), 5.8 (d, 0.5H), 6.1 (d, 0.5H), 7.2 (m, 6H), 8.55 (m, 1H), 11.85 (m, 1H); MS m/z 416, 418 (M-H).

25

The following example was made by the process of **Example 13**, using 2-chloro-N-[(1R,2R)-1-(methylamino)-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-b]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**) as the amine salt and the (S)-lactic acid as the carboxylic acid.


**Example 14: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2-hydroxypropanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



5 <sup>1</sup>H NMR (mixture of rotamers) δ: 1.1 (d, 1.5H), 1.22 (d, 1.5H), 2.64 (s, 1.5H), 2.83 (s, 1.5H), 3.0 (m, 1H), 3.2 (m, 1H), 4.5 (m, 1H), 4.8 (m, 2H), 5.7 (d, 0.5H), 6.15 (d, 0.5H), 7.2 (m, 6H), 8.55 (m, 1H), 11.85 (m, 1H); MS m/z 416, 418 (M-H).

**Example 15: 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2,3-**

10 **dihydroxypropanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**




2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[[[(4*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-

15 carboxamide (**Intermediate 20**; 285 mg, 0.56 mmol) was dissolved in acetic acid (4 mL, 20% aqueous) and warmed to 70°C for 3 h. The reaction was cooled, water (20 mL) added and the mixture filtered, the solid washed with water and dried *in vacuo* to give the title compound (200 mg, 76%) as a powder.

<sup>1</sup>H NMR (mixture of rotamers) δ: 2.65 (s, 1.5H), 2.9 (s, 1.5H), 3.0 (m, 1H), 3.25 (m, 1H), 3.5 (m, 2H), 4.65 (m, 4H), 5.65 (d, 0.5H), 6.1 (d, 0.5H), 7.05 (m, 2H), 7.25 (m, 3H), 8.63 (m, 1H), 12.4 (m, 1H); MS m/z 466, 468 (M-H).


The following example was made by the process of **Example 15**, using 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[[[(4*S*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide (**Intermediate 21**) as the acetone-

**Example 16: 2,3-Dichloro-*N*-{[(1*R*,2*R*)-1-[(2*S*)-2,3-dihydroxypropanoyl](methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



<sup>1</sup>H NMR (mixture of rotamers)  $\delta$ : 2.62 (s, 1.5H), 2.87 (s, 1.5H), 2.98 (m, 1H), 3.2 (m, 1H),  
 5 3.5 (m, 2H), 4.4 (m, 1.5H), 4.8 (m, 2H), 5.3 (m, 0.5H), 5.75 (d, 0.5H), 6.2 (d, 0.5H), 7.2 (m,  
 5H), 8.65 (d, 1H), 12.4 (m, 1H); MS *m/z* 466, 468 (M-H).

**Example 17: (2*S*)-*N*<sup>1</sup>-[(1*R*,2*R*)-2-[(2-Chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)-2-hydroxy-*N*<sup>1</sup>-methylsuccinamide**



10

(2*S*)-4-Amino-2-hydroxy-4-oxobutanoic acid (CAS Reg. No.: 57229-74-0, 109 mg, 0.82  
 mmol), 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-  
*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**; 284 mg, 0.74 mmol), HOBT (111  
 mg, 0.82 mmol) and triethylamine (0.46 mL, 3.3 mmol) were suspended in DMF (5 mL) and  
 15 stirred at ambient temperature. EDCI (157 mg, 0.82 mmol) was added and stirring was  
 continued at ambient temperature for a further 18 hours. The reaction mixture was purified by  
 reverse phase HPLC (5-95% acetonitrile / water gradient containing 0.2% TFA) to give the  
 title compound (34 mg, 10%) as a white solid.

<sup>1</sup>H NMR  $\delta$ : 2.43 (m, 2H), 2.75 (d, 3H), 2.97 (m, 1H), 3.22 (m, 1H), 4.77 (m, 2H), 5.90 (dd,  
 20 1H), 6.81 (d, 1H), 7.16 (m, 7H), 8.58 (dd, 1H), 11.85 (s, 1H); MS *m/z* 461.1.

The following examples were made by the process of **Example 17** using 2,3-dichloro-*N*-  
 [(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4,6a-dihydro-3a*H*-thieno[3,2-

*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**) as the amine source and the appropriate commercially available carboxylic acid.

**Example 18:** (2*S*)-*N*<sup>1</sup>-((1*R*,2*R*)-2-[[2,3-Dichloro-4*H*-thieno[3,2-*b*]pyrrol-5-

5 yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)-2-hydroxy-*N*<sup>1</sup>-methylsuccinamide

**Example 19:** 2,3-dichloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2-hydroxybutanoyl](methyl)amino}-2,3-  
dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide


**Example 20:** 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2-hydroxy-3-methylbutanoyl](methyl)  
amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide

10 **Example 21:** 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-4-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-  
yl)-2-hydroxybutanoyl](methyl)amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-  
*b*]pyrrole-5-carboxamide

**Example 22:** 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2-hydroxy-3-(methylthio)propanoyl]  
(methyl)amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide





15 **Example 23:** *tert*-Butyl {(2*S*)-3-[(1*R*,2*R*)-2-[[2,3-dichloro-4*H*-thieno[3,2-*b*]pyrrol-5-  
yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)(methyl)amino}-2-hydroxy-3-  
oxopropyl}carbamate

**Example 24:** 2,3-dichloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-3-cyano-2-hydroxypropanoyl](methyl)  
amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide




20

Example No	R Group	<sup>1</sup> H NMR δ:	MS
18		2.48 (m, 2H), 2.75 (d, 3H), 2.97 (m, 1H), 3.22 (m, 1H), 4.77 (m, 2H), 5.90 (dd, 1H), 6.81 (d, 1H), 7.14 (m, 6H), 8.67 (dd, 1H), 12.39 (s, 1H); MS m/z 493.2.	493.2
19		0.61 (t, 3H), 0.90 (t, 3H), 1.54 (m, 2H), 2.73 (d, 3H), 3.00 (m, 1H), 3.23 (m, 1H), 4.26 (m, 1H), 4.88 (m, 1H), 5.91 (dd, 1H), 6.98 (m, 2H), 7.26 (m, 3H), 8.69 (dd, 1H), 12.44 (d, 1H)	466.0

20		0.63 (dd, 3H), 0.87 (t, 3H), 1.94 (m, 1H), 2.73 (d, 3H), 3.00 (m, 1H), 3.23 (m, 1H), 4.05 (dd, 1H), 4.88 (m, 1H), 5.91 (dd, 1H), 6.98 (m, 2H), 7.26 (m, 3H), 8.67 (dd, 1H), 12.44 (d, 1H)	480.2
22		2.26 (s, 3H), 2.91 (m, 7H), 3.51 (m, 1H), 4.75 (m, 2H), 6.30 (d, 1H), 6.54 (s, 1H), 7.01 (d, 1H), 7.22 (m, 3H), 10.14 (s, 1H)	498.2
Intermediate 30		1.27 (d, 9H), 2.73 (d, 3H), 2.99 (m, 1H), 3.19 (m, 2H), 4.41 (m, 1H), 4.85 (m, 2H), 5.62 (dd, 1H), 6.14 (d, 1H), 7.08 (m, 5H), 8.63 (m, 1H), 12.40 (d, 1H)	564.5
24		2.90 (m, 4H), 3.07 (s, 3H), 3.41 (m, 1H), 4.88 (m, 2H), 6.25 (s, 1H), 6.37 (d, 1H), 6.68 (d, 1H), 7.03 (d, 3H), 7.23 (m, 2H), 10.58 (s, 1H)	475.1

There is no Example 21 or 23.

**Example 25: *N*-{(1*R*,2*R*)-1-[(*N*-Acetylseryl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-2-chloro-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



5

DIPEA (307  $\mu$ L, 1.8 mmol), HOBT (74 mg, 0.55 mmol), *N*-acetylserine (74 mg, 0.5 mmol) and EDAC (115 mg, 0.6 mmol) were added to a suspension of 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**; 191mg, 0.5 mmol) in anhydrous DMF (4 mL). The reaction was stirred at ambient temperature for approximately 16 h and then diluted with water (20 mL). The resulting crude product was recovered by filtration and purified by reverse phase

10

preparative HPLC (C18 ODS column, acetonitrile / water gradient 5-95% containing 0.2% TFA eluant) to give the title compound (35 mg, 8%).


$^1\text{H}$  NMR  $\delta$ : 1.8 (m, 3H), 3.0 (m, 7H), 4.9 (m, 2H), 5.7 (dd, 0.5H), 6.15 (dd, 0.5H), 7.1 (m, 6H), 8.25 (m, 2H), 11.8 (m, 1H); MS  $m/z$  473, 475.

5

The following compound was prepared by the process of **Example 25** using 2,3-dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**) as the amine and *N*-acetylserine as the carboxylic acid.


10

**Example 26 : *N*-{[(1*R*,2*R*)-1-[(*N*-Acetylseryl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl]-2,3-dichloro-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



$^1\text{H}$  NMR  $\delta$ : 1.75 (m, 3H), 3.0 (m, 7H), 4.8 (m, 2H), 5.7 (dd, 0.5H), 6.11 (dd, 0.5H), 7.2 (m, 5H), 8.3 (m, 2H), 12.35 (m, 1H); MS  $m/z$  507, 509.

**Example 27: 2,3-Dichloro-*N*-{[(1*R*,2*R*)-1-[methyl(*L*-seryl)amino]-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride**




DIPEA (266  $\mu\text{L}$ , 1.53 mmol), HOBt (101 mg, 0.75 mmol), *N*-(*tert*-butoxycarbonyl)-*L*-serine (103 mg, 0.5 mmol) and EDAC (119 mg, 0.62 mmol) were added to a suspension of 2,3-

20

dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**; 208 mg, 0.5 mmol) in anhydrous DMF (4 mL). The reaction was stirred at ambient temperature for approximately 16 h then diluted with water (20 mL) and the precipitated solid recovered by filtration and dried under vacuum. The crude material was purified by chromatography on silica gel eluting with an isohexane/EtOAc gradient (0-80%) and then dissolved in 4M HCl / Dioxan and left to stand for 1 h at ambient temperature. The volatiles were removed by evaporation under reduced pressure and the resulting gum triturated with ether to give the title compound (119 mg, 51%) as a white solid. <sup>1</sup>H NMR δ: 2.7 (s, 1.5H), 2.9 (s, 1.5H), 3.1 (m, 1H), 3.25 (m, 1H), 3.8 (m, 2H), 4.4 (m, 1H), 5.0 (m, 1H), 5.6 (m, 1H), 5.75 (d, 0.5H), 6.15 (d, 0.5H), 7.1 (m, 5H), 8.1 (m, 3H), 8.8 (d, 0.5H), 9.3 (d, 0.5H), 12.52 (d, 1H); MS m/z 467.

The following example was prepared by the method of **Example 27** using 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**) as the amine and *N*-(*tert*-butoxycarbonyl)-*L*-serine as the carboxylic acid.


**Example 28: 2-Chloro-*N*-{(1*R*,2*R*)-1-[methyl(*L*-seryl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride**



<sup>1</sup>H NMR δ: 2.7 (s, 1.5H), 2.9 (s, 1.5H), 3.1 (m, 1H), 3.25 (m, 1H), 3.8 (m, 2H), 4.4 (m, 1H), 5.0 (m, 1H), 5.6 (m, 1H), 5.8 (d, 0.5H), 6.2 (d, 0.5H), 7.25 (m, 6H), 8.1 (m, 3H), 8.7 (d, 0.5H), 9.1 (d, 0.5H), 11.9 (d, 1H); MS m/z 433.



**Example 29: (2S)-N<sup>1</sup>-((1R,2R)-2-[[2-Chloro-6H-thieno[2,3-b]pyrrol-5-yl]carbonyl]amino)-2,3-dihydro-1H-inden-1-yl)-2-hydroxy-N<sup>1</sup>-methylpentanediamide**




2-Chloro-N-[(1R,2R)-1-(methyl{[(2S)-5-oxotetrahydrofuran-2-yl]carbonyl}amino)-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-b]pyrrole-5-carboxamide (**Intermediate 22**; 100 mg, 0.22 mmol) was suspended in ammonia (5 mL, 2M in isopropanol, 2.5 mmol) and the mixture heated by microwave irradiation at 150 °C for 30 min. After evaporation of the reaction mixture the crude product was purified by reverse phase preparative HPLC (C18 ODS column, acetonitrile / water gradient 5-95% containing 0.2% TFA eluant) to give the title compound (52 mg, 49%).

<sup>1</sup>H NMR δ: 1.65 (m, 1H), 2.0 (m, 1H), 2.2 (m, 2.0H), 2.65 (s, 1.5H), 2.9 (s, 1.5H), 3.0 (m, 1H), 3.3 (m, 1H), 4.35 (m, 1H), 4.7 (d, 0.5H), 4.9 (m, 1H), 5.3 (d, 0.5H), 5.7 (d, 0.5H), 6.2 (d, 0.5H), 6.8 (d, 1H), 7.0 (m, 1.5H), 7.1 (m, 0.5H), 7.2 (s, 1H), 7.35 (m, 4H), 8.65 (m, 1H), 11.9 (s, 1H); MS m/z 475.

The following example was prepared by the method of **Example 29** using 2,3-dichloro-N-[(1R,2R)-1-(methyl{[(2S)-5-oxotetrahydrofuran-2-yl]carbonyl}amino)-2,3-dihydro-1H-inden-2-yl]-4H-thieno[3,2-b]pyrrole-5-carboxamide (**Intermediate 23**) as the lactone.


**Example 30: (2S)-N<sup>1</sup>-((1R,2R)-2-[[2,3-Dichloro-4H-thieno[3,2-b]pyrrol-5-yl]carbonyl]amino)-2,3-dihydro-1H-inden-1-yl)-2-hydroxy-N<sup>1</sup>-methylpentanediamide**



<sup>1</sup>H NMR δ: 1.6 (m, 1H), 2.1 (m, 2H), 2.45 (m, 1H), 2.65 (s, 1.5H), 2.85 (s, 1.5H), 3.0 (m, 1H), 3.3 (m, 1H), 4.3 (m, 1H), 4.65 (d, 0.5H), 4.9 (m, 1H), 5.25 (d, 0.5H), 5.7 (d, 0.5H), 6.1 (d, 0.5H), 6.7 (m, 1H), 6.9 (m, 0.5H), 7.1 (m, 1.5H), 7.25 (m, 4H), 8.7 (m, 1H), 12.4 (s, 1H); MS m/z 509.

5

**Example 31: 2-Chloro-*N*-{[(1*R*,2*R*)-1-[(2*S*)-2-hydroxy-3-methoxypropanoyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl}-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**




2-Chloro-*N*-((1*R*,2*R*)-1-{methyl[(2*S*)-oxiran-2-ylcarbonyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide (**Intermediate 24**; 100 mg, 0.24 mmol) was suspended in sodium methoxide (5 mL, 0.5M solution in methanol) and heated under microwave irradiation at 100 °C for 5 min. Acetic acid (0.5 mL) was then added and the reaction mixture evaporated. The residue was then purified by reverse phase preparative HPLC (C18 ODS column, acetonitrile / water gradient 5-95% containing 0.2% TFA eluant) to give the title compound (33 mg, 32%).

<sup>1</sup>H NMR δ: 2.6 (s, 1.5H), 2.8 (s, 1.5H), 3.0 (m, 1H), 3.05 (s, 1.5H), 3.2 (m, 1H), 3.3 (s, 1.5H), 3.45 (m, 2H), 4.5 (t, 1H), 4.85 (m, 1H), 5.7 (d, 0.5H), 6.15 (d, 0.5H), 7.0 (m, 1.5H), 7.1 (m, 0.5H), 7.2 (s, 1H), 7.25 (m, 3H), 8.6 (t, 1H), 11.86 (d, 1H); MS m/z 448.


The following example was prepared by the method of **Example 31** using 2,3-dichloro-*N*-((1*R*,2*R*)-1-{methyl[(2*S*)-oxiran-2-ylcarbonyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide (**Intermediate 25**) as the oxirane.

**Example 32: 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2-hydroxy-3-methoxypropanoyl](methyl)amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



5 <sup>1</sup>H NMR δ: 2.6 (s, 1.5H), 2.8 (s, 1.5H), 3.0 (m, 1H), 3.05 (s, 1.5H), 3.2 (m, 1H), 3.3 (s, 1.5H), 3.45 (m, 2H), 4.55 (t, 1H), 4.9 (m, 1H), 5.75 (d, 0.5H), 6.2 (d, 0.5H), 6.95 (m, 0.5H), 7.1 (m, 1.5H), 7.3 (m, 3H), 8.7 (t, 1H), 12.42 (d, 1H); MS m/z 482.


**Example 33: 2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2-hydroxy-3-(methylsulfonyl)propanoyl](methyl)amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



2,3-Dichloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2-hydroxy-3-(methylthio)propanoyl](methyl)amino}-2,3-dihydro-1*H*-inden-2-yl}-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide (**Example 22**; 200 mg, 0.40 mmol) was dissolved in DMF (5 mL) at 0 °C, and *m*-CPBA (218 mg, 70% strength, 0.88 mmol) was added. The reaction was stirred for 3 h at ambient temperature before purification by reverse phase HPLC (C18 ODS column, 5-95% acetonitrile / water gradient containing 0.2% TFA) to give the title compound (150 mg, 70%) as a white solid.

<sup>1</sup>H NMR δ: 2.80 (d, 3H), 2.89 (d, 3H), 2.97 (m, 1H), 3.22 (m, 1H), 3.41 (dd, 2H), 4.88 (m, 2H), 5.90 (dd, 1H), 7.18 (m, 5H), 8.70 (dd, 1H), 12.39 (d, 1H); MS m/z 530.1.

**Example 34: *N*-{[(1*R*,2*R*)-1-[(2*S*)-3-Amino-2-hydroxypropanoyl](methylamino)-2,3-dihydro-1*H*-inden-2-yl]-2,3-dichloro-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride**



5 *tert*-Butyl {(2*S*)-3-[[[(1*R*,2*R*)-2-[[2,3-dichloro-4*H*-thieno[3,2-*b*]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1*H*-inden-1-yl](methylamino)-2-hydroxy-3-oxopropyl]carbamate


(**Intermediate 30**; 291 mg, 0.51 mmol) was dissolved in DCM (15 mL), 4M HCl/ dioxane (15 mL) was added and the reaction mixture stirred for 1 h. The solvent was evaporated, the residue slurried in ether (30 mL) and filtered to give the title compound (190 mg, 74%) as a

10 white solid.

<sup>1</sup>H NMR δ: 2.77 (d, 3H), 3.03 (m, 2H), 3.25 (m, 1H), 4.76 (m, 2H), 5.79 (d, 1H), 6.08 (d, 1H), 7.18 (m, 5H), 7.90 (s, 3H), 8.90 (dd, 1H), 12.48 (d, 1H); MS *m/z* 466.9.

**Example 35: (2*S*)-*N*<sup>1</sup>-[(1*R*,2*R*)-2-[[2,3-Dichloro-4*H*-thieno[3,2-*b*]pyrrol-5-**

15 **yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)-2-hydroxy-*N*<sup>1</sup>,*N*<sup>4</sup>-dimethylsuccinamide**



Methylamine (11 mL, 2M in THF, 22 mmol), HOBt (1.49 g, 11 mmol), [(4*S*)-2,2-dimethyl-5-oxo-1,3-dioxolan-4-yl]acetic acid (1.74 g, 10 mmol) and EDAC (2.11 g, 11 mmol) were dissolved in DCM (50 mL). The reaction was stirred at ambient temperature for

20 approximately 16 h before evaporation and purification by flash column chromatography on silica gel eluting with 6-10% MeOH/ DCM. The resulting gum was dissolved in 4M HCl/ dioxane (30 ml) before adding water (1.5 ml) and stirring for 1 h. The solvent was evaporated to afford the crude acid. This acid (448 mg, 3.0 mmol), 2,3-dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4,6a-dihydro-3a*H*-thieno[3,2-*b*]pyrrole-5-


25 carboxamide hydrochloride (**Intermediate 16**; 423 mg, 1.0 mmol), HOBt (205 mg, 1.5

mmol), and triethylamine (0.62 ml, 4.5 mmol) were suspended in DMF (15 mL) and stirred at room temperature. EDCI (292 mg, 1.5 mmol) was added and stirring was continued for a further 18 hours. The reaction mixture was purified by reverse phase HPLC (C18 ODS column 5-95% acetonitrile / water gradient containing 0.2% TFA) to give the title compound 5 (30 mg, 6%) as a white solid.

$^1\text{H}$  NMR  $\delta$ : 2.71 (m, 9H), 3.50 (m, 1H), 4.91 (m, 2H), 5.95 (dd, 1H), 6.85 (m, 2H), 7.24 (m, 5H), 8.10 (d, 1H), 10.36 (d, 1H); MS  $m/z$  508.9.


The following example was made by the process of **Example 35** using dimethylamine as the amine.

**Example 36: (2S)-N<sup>1</sup>-((1R,2R)-2-[[2,3-Dichloro-4H-thieno[3,2-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1H-inden-1-yl)-2-hydroxy-N<sup>1</sup>,N<sup>4</sup>,N<sup>4</sup>-trimethylsuccinamide**



$^1\text{H}$  NMR  $\delta$ : 2.93 (m, 9H), 3.49 (s, 3H), 3.58 (m, 1H), 4.99 (m, 2H), 6.03 (dd, 1H), 7.00 (m, 7H), 8.39 (d, 1H), 10.17 (s, 1H); MS  $m/z$  522.9.

**Example 37: 2-Chloro-N-((1R,2R)-1-[[glyceroyl(2-hydroxyethyl)amino]-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-*b*]pyrrole-5-carboxamide**



20


2-Chloro-N-((1R,2R)-1-[[2,2-dimethyl-1,3-dioxolan-4-yl)carbonyl][2-(tetrahydro-2H-pyran-2-yloxy)ethyl]amino}-2,3-dihydro-1H-inden-2-yl)-6H-thieno[2,3-*b*]pyrrole-5-carboxamide (**Intermediate 26**; 250 mg, 0.472 mmol) was dissolved in acetic acid (6 mL, 20% aqueous) and warmed to 60°C for 4h. The reaction was cooled and the volatiles removed under reduced

pressure. The residue was dissolved in EtOAc (15 mL), washed with saturated NaHCO<sub>3</sub> (5 mL), water (5 mL) and brine (5 mL), dried (MgSO<sub>4</sub>) and the volatiles removed under reduced pressure. The residue was purified by flash column chromatography (EtOAc) to give the title compound (110mg, 56%) as a foam.


- 5 <sup>1</sup>H NMR (mixture of rotamers and diastereomers)  $\delta$ : 2.8(m, 2H), 3.4(m, 6H), 4.8(m, 5H), 5.68(d, 0.5H), 5.8(d, 0.5H), 7.2(m, 6H), 8.55(m, 1H), 11.9(s, 1H); MS m/z 462, 464 (M-H).

The following examples were made by the process of **Example 37**, using 2-Chloro-*N*-((1*R*,2*R*)-1-{[(*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl}[2-(tetrahydro-2*H*-pyran-2-yloxy)ethyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide (**Intermediate 27**) or 2-chloro-*N*-((1*R*,2*R*)-1-{[(*S*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl}[2-(tetrahydro-2*H*-pyran-2-yloxy)ethyl]amino)-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide (**Intermediate 28**) as the protected triol.


- 15 **Example 38: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(2*R*)-2,3-dihydroxypropanoyl](2-hydroxyethyl)amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**  
**Example 39: 2-Chloro-*N*-{(1*R*,2*R*)-1-[(2*S*)-2,3-dihydroxypropanoyl](2-hydroxyethyl)amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**




Example	R	<sup>1</sup> H NMR $\delta$	M/z
38		2.8(m, 2H), 3.4(m, 6H), 4.8(m, 5H), 5.68(d, 1H), 7.2(m, 6H), 8.55(m, 1H), 11.9(s, 1H)	462, 464 (M-H)
39		2.8(m, 2H), 3.4(m, 6H), 4.8(m, 5H), 5.8(d, 1H), 7.2(m, 6H), 8.55(m, 1H), 11.9(s, 1H)	462, 464 (M-H)

**Intermediates:****Intermediate 1: 2-Chloro-N-[(1R,2R)-1-(methylamino)-2,3-dihydro-1H-inden-2-yl]-6H-thieno[2,3-b]pyrrole-5-carboxamide hydrochloride salt**


- 5 *tert*-Butyl ((1R,2R)-2-[[[(2-chloro-6H-thieno[2,3-b]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1H-inden-1-yl)methylcarbamate (**Intermediate 2**; 5.8g, 13.0mmol) was dissolved in HCL solution (4N in dioxane, 40ml) and stirred at ambient temperature for 24 hours. The volatiles were removed by evaporation under reduced pressure and the residue dried *in vacuo* to give the title compound (3.0g, 60%) as a powder.
- 10 <sup>1</sup>H NMR δ: 2.7(s, 3H), 3.05(dd, 1H), 3.5(dd, 1H), 4.82(m, 2H), 7.09(s, 1H), 7.15(s, 1H), 7.35(m, 3H), 7.78(d, 1H), 8.9(d, 1H), 9.55(broad d, 2H), 11.95(s, 1H); MS m/z 346, 348.

**Intermediate 2: *tert*-Butyl ((1R,2R)-2-[[[(2-chloro-6H-thieno[2,3-b]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1H-inden-1-yl)methylcarbamate**

- 15 2-Chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid (**Intermediate 3**; 3.5 g, 17.16 mmol), *tert*-butyl [(1R,2R)-2-amino-2,3-dihydro-1H-inden-1-yl)methylcarbamate (**Intermediate 5**; 4.5 g, 17.16 mmol), DIPEA (2.9 ml, 17.16 mmol) and HOBt (2.32 g, 17.16 mmol) were dissolved in DCM (60 ml), stirred for 5 minutes, EDCI (4.11 g, 21.45 mmol) added and the
- 20 mixture stirred at ambient temperature for 24 hours. The volatiles were removed by evaporation under reduced pressure and EtOAc (200 mL) added. The organic phase was washed with water (100 mL), brine (50 mL) and dried (MgSO<sub>4</sub>), filtered and the solvent removed under reduced pressure. The residue was purified by column chromatography (SiO<sub>2</sub>, EtOAc:Hexane) to afford the title compound (5.8g, 76%) as an oil.
- 25 <sup>1</sup>H NMR δ: 1.3 (s, 9H), 2.63 (s, 3H), 3.1 (m, 2H), 4.75 (m, 1H), 5.63 (m, 1H), 7.0 (m, 2H), 7.16 (s, 1H), 7.25 (m, 3H), 8.48 (m, 1H), 11.83 (s, 1H).

**Intermediate 3: 2-Chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid**


- Sodium hydroxide (2N, 15 mL) was added to a methanol (50 mL) solution of 2-chloro-5-methoxycarbonyl-6H-thieno[2,3-b]pyrrole (**Intermediate 4**, 777 mg, 3.6 mmol) and the mixture heated at reflux for 5 hours. The reaction was cooled to ambient temperature, water (250 mL) added and the aqueous phase was washed with Et<sub>2</sub>O (2 x 50 mL), acidified to pH 2 with HCl (2N) and extracted with EtOAc (3 x 50 mL). The combined organic phases were washed with water (2 x 50 mL), brine (50 mL), dried (MgSO<sub>4</sub>) and the solvent removed under reduced pressure to afford the title compound (705 mg, 97%) as a pale pink solid.
- <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 12.6-12.7 (1H, b), 12.0-12.1 (1H, b), 7.15 (1H, s), 6.9 (1H, s); m/z 183, 185.

**Intermediate 4: 2-Chloro-5-methoxycarbonyl-6H-thieno[2,3-b]pyrrole**

- Sodium (659 mg, 28.7 mL) was added to dry methanol (20 mL) and the mixture stirred at ambient temperature for 30 mins before cooling to -20 °C. 2-Chlorothiophene-3-carboxaldehyde (Gronowitz *et al.*, Tetrahedron Vol.32 1976 p.1403; 1.17 g, 7.2 mmol) and methyl azidoacetate (3.3 g, 28.7 mmol) were added as a methanol (10 mL) solution and the reaction was stirred from -20 °C to 10 °C over 16 hours. The reaction was poured on saturated NH<sub>4</sub>Cl (300 mL) and extracted with DCM (3 x 100 mL). The combined organic phases were washed with water (2 x 100 mL), brine (100 mL), dried (MgSO<sub>4</sub>) and the solvent removed under reduced pressure. The crude product was redissolved in xylene (50 mL) and added dropwise to refluxing xylene (150 mL) and stirred for at reflux for a further 30 mins after the addition was complete. The solvent was removed under reduced pressure to afford a yellow solid which was recrystallised (25:75, EtOAc:isohexane) to afford the title compound (1.06 g, 69%) as a solid.
- <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 9.4-9.2 (1H, br), 7.0 (1H, s), 6.9 (1H, s), 3.9 (3H, s); m/z 214, 216.



**Intermediate 5: *tert*-Butyl [(1*R*,2*R*)-2-amino-2,3-dihydro-1*H*-inden-1-yl]methyl carbamate**




(1*R*,2*S*)-1-[(*tert*-Butoxycarbonyl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl

- 5 methanesulfonate (**Intermediate 6**; 3.0g, 8.8mmol) and sodium azide (2.3 g, 35.2 mmol) in dry DMA (30 mL) was heated to 90°C for 7 hours. The reaction was cooled and ethyl acetate (100 mL) added. The mixture was washed with water (6 x 25 mL), brine (50 mL) and dried (MgSO<sub>4</sub>). 10% Palladium on carbon (400 mg) was added to the organic solution which was stirred under a hydrogen atmosphere for 4h, filtered through Celite and evaporated. The
- 10 residue was purified by column chromatography (EtOAc and then DCM:MeOH 9:1) to afford the title compound (1.2 g, 55%) as a pale brown oil.

<sup>1</sup>H NMR  $\delta$ : 1.45 (m, 9H), 2.6 (s, 3H), 2.8 (m, 1H), 3.3 (m, 1H), 4.45 (m, 1H), 5.55 (dd, 1H), 7.26 (m, 4H); MS *m/z* 264.


- 15 **Intermediate 6: (1*R*,2*S*)-1-[(*tert*-Butoxycarbonyl)(methyl)amino]-2,3-dihydro-1*H*-inden-2-yl methanesulfonate**



- tert*-Butyl [(1*R*,2*S*)-2-hydroxy-2,3-dihydro-1*H*-inden-1-yl]methylcarbamate (**Intermediate 7**; 3.0 g, 11.4 mmol) was dissolved in dry THF (40 mL) at 10°C. A solution of methane
- 20 sulphonyl chloride (1.44 g, 12.55 mmol) in dry THF (10 mL) was added, the reaction allowed to warm to ambient temperature and stirred for 30 mins. The volatiles were removed by evaporation under reduced pressure and ethyl acetate (100 mL) added. The mixture was washed with water (2 x 50 mL), brine (50 mL) and the organic phase was dried (MgSO<sub>4</sub>), filtered and evaporated. The residue was purified by column chromatography
- 25 (EtOAc:Hexane) to afford the title compound (3.1g, 80%) as a colourless syrup.

<sup>1</sup>H NMR  $\delta$ : 1.46 (s, 9H), 2.61 (s, 3H), 3.12 (m, 1H), 3.18 (s, 3H), 3.32 (m, 1H), 5.45 (m, 1H), 5.68 (m, 1H), 7.28 (m, 4H); MS *m/z* 342.


**Intermediate 7: *tert*-Butyl [(1*R*,2*S*)-2-hydroxy-2,3-dihydro-1*H*-inden-1-yl]methylcarbamate**



*tert*-Butyl methyl[(1*R*,2*S*)-2-(tetrahydro-2*H*-pyran-2-yloxy)-2,3-dihydro-1*H*-inden-1-yl]carbamate (**Intermediate 8**; 4.0 g, 11.5 mmol) was dissolved in methanol (50 mL), 4-toluene sulphonic acid added and the reaction stirred at ambient temperature for 2 hours. Saturated NaHCO<sub>3</sub> (50 mL), water (100 mL) was added and ethyl acetate (100 mL) was added and the mixture stirred for 30 mins. The organic phase was separated, washed with water (50 mL), brine (50 mL) and dried (MgSO<sub>4</sub>). The volatiles were removed by evaporation under reduced pressure to give the title compound (3.0 g, 99%) as an oil.

<sup>1</sup>H NMR  $\delta$ : 1.45 (s, 9H), 2.6 (s, 3H), 2.75 (m, 1H), 3.05 (m, 1H), 4.5 (m, 1H), 5.05 (m, 1H), 5.34 (m, 1H), 7.03-7.3 (m, 4H).


**Intermediate 8: *tert*-Butyl methyl[(1*R*,2*S*)-2-(tetrahydro-2*H*-pyran-2-yloxy)-2,3-dihydro-1*H*-inden-1-yl]carbamate**



*tert*-Butyl [(1*R*,2*S*)-2-(tetrahydro-2*H*-pyran-2-yloxy)-2,3-dihydro-1*H*-inden-1-yl]carbamate (**Intermediate 9**; 4.0 g, 12.0 mmol) was dissolved in dry DMA (25 mL) at 5°C. 60% Sodium hydride (575 mg, 14.4 mmol) was added, the reaction stirred at 5°C for 30 mins, allowed to warm to ambient temperature and stirred for a further 30 mins. Methyl iodide (896  $\mu$ L, 14.4 mmol) was added and the reaction stirred at ambient temperature for 3 hours. The reaction was poured into water (100 mL) and extracted with ethyl acetate (2 x 50ml). The organic extracts were washed with water (6 x 25 mL), brine (50 mL) and dried (MgSO<sub>4</sub>). The volatiles were removed by evaporation under reduced pressure to give the title compound (4.1 g, 97%) as an oil.

<sup>1</sup>H NMR  $\delta$ : 1.4-1.9 (m, 6H), 1.5 (s, 9H), 2.7 (dd, 3H), 2.85-3.3 (m, 2H), 3.5 (m, 1H), 3.7-4.0 (m, 1H), 4.6-4.9 (m, 2H), 5.5-5.85 (m, 1H), 7.2 (s, 4H).

**Intermediate 9: *tert*-Butyl [(1*R*,2*S*)-2-(tetrahydro-2*H*-pyran-2-yl)oxy]-2,3-dihydro-1*H*-inden-1-yl]carbamate**




*tert*-Butyl [(1*R*,2*S*)-2-hydroxy-2,3-dihydro-1*H*-inden-1-yl]carbamate (**Intermediate 10**, 7.0 g, 28.1 mmol) and 3,4-dihydro-2*H*-pyran (4.7 g, 56.2 mmol) dissolved in DCM (50 mL). 4-Toluenesulphonic acid pyridinium salt (100 mg) was added and the reaction stirred for 4 hours at ambient temperature. The reaction was diluted with ethyl acetate (100 mL), washed with water (2 x 50 mL), brine (50 mL) and dried (MgSO<sub>4</sub>). The volatiles were removed by evaporation under reduced pressure to give the title compound (8.9 g, 95%) as an oil.

<sup>1</sup>H NMR  $\delta$ : 1.25-1.85 (m, 6H), 1.45 (s, 9H), 2.85-3.1 (m, 2H), 3.4 (m, 1H), 3.8 (m, 1H), 4.35-5.1 (m, 3H), 6.8 (dd, 1H), 7.2(s, 1H).

15


**Intermediate 10: *tert*-Butyl [(1*R*,2*S*)-2-hydroxy-2,3-dihydro-1*H*-inden-1-yl]carbamate**



(1*R*,2*S*)-1-Amino-2,3-dihydro-1*H*-inden-2-ol (CAS Reg. No. 136030-00-7; 10 g, 67.1 mmol) was dissolved in DCM (550 mL) and Et<sub>3</sub>N (18.7 mL, 134.2 mmol). Di-*tert*-butyl dicarbonate (18.3 g, 83.9 mmol) in DCM (50 mL) was added and the mixture stirred at ambient temperature for 20 hours, and then evaporated. EtOAc (200 mL) was added, the solution washed with water (200 mL), dried (MgSO<sub>4</sub>) and the volatiles removed under reduced pressure. The crude product was purified by flash column chromatography (SiO<sub>2</sub>, 4:1, *iso*-hexane:EtOAc eluent) to provide the title compound (16.1 g, 96%) as a white solid.

<sup>1</sup>H NMR  $\delta$ : 1.42 (m, 9H), 2.78 (dd, 1H), 3.00 (dd, 1H), 4.36 (m, 1H), 4.84 (m, 1H), 4.95 (m, 1H), 6.3 (d, 1H), 7.13 (m, 4H).


**Intermediate 11: Ethyl *N*-[(acetyloxy)acetyl]-*N*-((1*R*,2*R*)-2-[[2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)glycinate**



Ethyl *N*-((1*R*,2*R*)-2-[[2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)glycinate (**Intermediate 12**; 820 mg, 2.0 mmol), acetoxyacetic acid (236 mg, 2.0 mmol) and HOBT (270 mg, 2.0 mmol) were dissolved in DCM (20ml), stirred for 5 minutes, EDCI (480 mg, 2.5 mmol) added and the mixture stirred at ambient temperature for 4 hours. The volatiles were removed by evaporation under reduced pressure and EtOAc (25 mL) added. The mixture was washed with water (2 x 10 mL), brine (10 mL) and dried (MgSO<sub>4</sub>), filtered and solvent removed under reduced pressure. The residue was purified by column chromatography (SiO<sub>2</sub>, EtOAc:Hexane) to afford the title compound (750mg, 75%) as a foam.

<sup>1</sup>H NMR  $\delta$ : 1.1(t, 3H), 2.02(s, 1.5H), 2.08(s, 1.5H), 2.9(m, 1H), 3.25(m, 1H), 3.68(m, 1H), 4.0(m, 3H), 4.68(m, 1H), 5.0(m, 2H), 5.45(d, 0.7H), 5.9(d, 0.3H), 7.2(m, 6H), 8.45(d, 0.3H), 8.55(d, 0.7H), 11.8(s, 0.3H), 11.84(s, 0.7H).

**Intermediate 12: Ethyl *N*-((1*R*,2*R*)-2-[[2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)glycinate**




*N*-[(1*R*,2*R*)-1-Amino-2,3-dihydro-1*H*-inden-2-yl]-2-chloro-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride salt (**Intermediate 13**, 1.8 g, 5.0 mmol), DIPEA (856  $\mu$ l, 5.0

mmol), ethyl glyoxalate (1.02 ml, 50% solution in toluene, 5.0 mmol) and acetic acid (300  $\mu$ l, 5.0 mmol) were dissolved in anhydrous THF (40 ml) and stirred at ambient temperature for 1 hour. Sodium triacetoxo borohydride (1.6 g, 7.5 mmol) was added, the reaction stirred at ambient temperature for 4 hours. HCl (1N, 10ml) and EtOAc (75ml) were added and the phases separated. The organic phase was washed with saturated NaHCO<sub>3</sub> (25ml), water (25ml), brine (25ml), dried (MgSO<sub>4</sub>) filtered and the solvent removed under reduced pressure. The residue was purified by column chromatography (SiO<sub>2</sub>; EtOAc:Hexane) to afford the title compound (1.7 g, 81%) as a solid.

<sup>1</sup>H NMR  $\delta$ : 1.27 (t, 3H), 1.80 (br s, 1H), 2.82 (dd, 1H), 3.7 (m, 3H), 4.22 (m, 3H), 4.5 (m, 1H), 6.78 (m, 2H), 6.85 (s, 1H), 7.25 (m, 3H), 7.37 (m, 1H), 10.68 (s, 1H).


**Intermediate 13: N-[(1R,2R)-1-Amino-2,3-dihydro-1H-inden-2-yl]-2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxamide hydrochloride salt**



tert-Butyl ((1R,2R)-2-[[[(2-chloro-6H-thieno[2,3-b]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1H-inden-1-yl]carbamate (**Intermediate 14**; 3.5 g, 8.1 mmol) was suspended in DCM (10 mL). HCl (20 mL, 4N in dioxane, 80 mmol) was added and reaction stirred at ambient temperature for 48 h. The volatiles were removed under reduced pressure; the resulting solid was azeotroped with EtOAc (4 x 50 mL) and dried *in vacuo* to afford the title compound (2.91 g, 97%) as a brown solid.

<sup>1</sup>H NMR  $\delta$ : 3.04 (dd, 1H), 3.87 (dd, 1H), 4.65 (m, 1H), 4.83 (br s, 1H), 7.15 (m, 2H), 7.33 (m, 3H), 7.69 (d, 1H), 8.80 (br s, 3H), 8.96 (d, 1H), 12.00 (s, 1H). m/z 330, 332 [M-H]<sup>+</sup>.


**Intermediate 14: *tert*-Butyl ((1*R*,2*R*)-2-[[*(*2-chloro-6*H*-thieno[2,3-*b*]pyrrol-5-yl)carbonyl]amino}-2,3-dihydro-1*H*-inden-1-yl)carbamate**



(1*R*, 2*R*)-2-Amino-1-[(1,1-dimethylethoxy)carbonylamino]indan (**Intermediate 15**; 3.7 g, 15 mmol), 2-chloro-6*H*-thieno[2,3-*b*]pyrrole-5-carboxylic acid (**Intermediate 3**; 3.0 g, 15 mmol), HOBT (2.03 g, 15 mmol) and DIPEA (2.83 mL, 16.5 mmol) were suspended in DCM (60 mL) and stirred at ambient temperature for 5 mins. EDAC (3.57 g, 18.75 mmol) was added and the mixture stirred at ambient temperature for 20 hours. The resulting precipitate was filtered, washed with DCM (2 x 20 mL) and dried to afford the title compound (5.35 g, 10 67%) as a yellow solid.

<sup>1</sup>H NMR  $\delta$ : 1.38 (s, 9H), 2.81 (dd, 1H), 3.17 (dd, 1H), 4.56 (m, 1H), 5.14 (m, 1H), 7.01 (s, 1H), 7.16 (m, 5H), 7.32 (d, 1H), 8.47 (d, 1H), 11.82 (s, 1H).

**Intermediate 15: (1*R*, 2*R*)-2-Amino-1-[(1,1-dimethylethoxy)carbonylamino]indan**



15


*Cis*-1-[(1,1-Dimethylethoxy)carbonylamino]-2-hydroxyindan (**Intermediate 10**; 14.0g, 56.2mmol) was dissolved in DCM (200 ml) and triethylamine (11.8 ml, 84.3mmol). Methanesulfonyl chloride (7.1g, 61.9mmol) dissolved in DCM (20 ml) was added and the mixture stirred at room temperature for 3 hours. The mixture was evaporated and EtOAc (250 20 ml) added. After washing with water and drying over magnesium sulphate the organic solution was evaporated to yield *cis*-1-[(1,1-dimethylethoxy)carbonylamino]-2-methanesulphonyloxyindan (9.7g, 98%) as a white solid.

<sup>1</sup>H NMR  $\delta$ : 1.45 (s, 9H), 3.15 (m, 2H), 3.18 (s, 3H), 5.20 (m, 1H), 5.35 (m, 1H), 7.15 (m, 4H), 7.45 (d, 1H).

*Cis*-1-[(1,1-dimethylethoxy)carbonylamino]-2-methanesulphonyloxindan (18.1g, 55.3mmol) was dissolved in dry dimethyl acetamide (100 ml). Sodium azide (5.4g, 83.0mmol) was added and the mixture heated to 90°C for 6 hours. The reaction was cooled, diluted with ethyl acetate (150 ml), washed with water (6 x 200 ml) and dried over magnesium sulphate. 10% Palladium on activated carbon was added and the mixture stirred under a hydrogen atmosphere for 24 hours. Filtration through celite followed by evaporation gave the title compound (2.6g, 98%) as a white solid.

<sup>1</sup>H NMR: 1.45 (s, 9H), 2.50 (dd, 1H), 3.05 (dd, 1H), 3.30 (m, 3H), 4.55 (m, 1H), 7.1 (m, 5H).


10 **Intermediate 16: 2,3-Dichloro-N-[(1R,2R)-1-(methylamino)-2,3-dihydro-1H-inden-2-yl]-4,6a-dihydro-3aH-thieno[3,2-b]pyrrole-5-carboxamide hydrochloride**



This intermediate was prepared by the method of **Intermediate 1**, using: *tert*-butyl ((1R,2R)-2-[[[(2,3-dichloro-4,6a-dihydro-3aH-thieno[3,2-b]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1H-inden-1-yl)methylcarbamate (**Intermediate 17**) as the carbamate.


<sup>1</sup>H NMR δ: 2.69 (m, 3H), 3.04 (dd, 1H), 3.51 (m, 1H), 4.85 (m, 2H), 7.19 (s, 2H), 7.35 (m, 3H), 7.75 (d, 1H), 9.10 (d, 1H) 9.55 (d, 2H), 12.58 (s, 1H); MS m/z 378.2.

20 **Intermediate 17: *tert*-butyl ((1R,2R)-2-[[[(2,3-dichloro-4,6a-dihydro-3aH-thieno[3,2-b]pyrrol-5-yl)carbonyl]amino]-2,3-dihydro-1H-inden-1-yl)methylcarbamate**




This intermediate was made by the process of **Intermediate 2**, using *tert*-Butyl [(1R,2R)-2-amino-2,3-dihydro-1H-inden-1-yl)methylcarbamate (**Intermediate 5**) as the amine and 5-Carboxy-2,3-dichloro-4H-thieno[3,2-b]pyrrole (**Intermediate 18**) as the carboxylic acid.

25 <sup>1</sup>H NMR δ: 1.28 (d, 9H), 2.65 (s, 3H), 3.09 (m, 2H), 4.79 (m, 1H), 5.62 (dd, 1H), 7.05 (m, 2H), 7.24 (m, 3H), 8.59 (m, 1H), 12.42 (d, 1H); MS m/z 478.2.

**Intermediate 18: 5-Carboxy-2,3-dichloro-4H-thieno[3,2-b]pyrrole**

2,3-Dichloro-5-methoxycarbonyl-4H-thieno[3,2-b]pyrrole (**Intermediate 19**; 1.22 g, 5.66 mmol) was taken up in MeOH (10 mL) and was heated under reflux. Aqueous lithium hydroxide (3.0 mL, 2.0 M, 6.0 mmol) was added portionwise over 45 mins. The mixture was heated at reflux for 1 h before being cooled and concentrated. Water (15 mL) was added and the solution was neutralised using 2 M aqueous HCl (5.66 mL). The solution was extracted using EtOAc (3x20 mL), and the combined organic layers were concentrated to afford the title compound (1.18 g, 100%) as a yellow solid

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 7.0 (1H, s); MS m/z 234.


**Intermediate 19: 2,3-Dichloro-5-methoxycarbonyl-4H-thieno[3,2-b]pyrrole**

Sodium methoxide solution (5 mL, 28% w/v solution, 25.9 mmol) was diluted with MeOH (10 mL) and was cooled to -25 °C under nitrogen. A solution of 3,4-dichloro-5-carbonyl-4H-thieno[3,2,b]pyrrole (DE 2814798; 1.2 g, 6.63 mmol) and methyl azidoacetate (3.0 g, 26.09 mmol) in MeOH (15 mL) was added dropwise, maintaining the temperature at -25 °C. On completion of addition the solution was allowed to warm to 5 °C over a period of approximately 16 h. The solution was added to saturated aqueous ammonium chloride (250 mL) and the mixture was extracted using DCM. The combined organic layers were concentrated at ambient temperature. The residue was taken up in xylene (30 mL) and this solution was added dropwise to xylene (120 mL) under reflux. The solution was heated under reflux for 30 mins before being cooled and concentrated. The title compound was purified by crystallisation (EtOAc/isohexane) to afford the title compound (1.08 g, 65%) as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 9.2 (1H, br), 7.0 (1H, s), 3.9 (3H, s); MS m/z 248.



**Intermediate 20: 2,3-Dichloro-*N*-[(1*R*,2*R*)-1-[[*(4R*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl](methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**




5 2,3-Dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**; 320mg, 0.768 mmol), (*4R*)-2,2-dimethyl-1,3-dioxolane-4-carboxylic acid potassium salt (212mg, 1.15 mmol) and DIPEA (131μL, 0.768mmol) in anhydrous DMA (5ml) was added HOBT (104mg, 0.768 mmol) and EDCI (184mg, 0.96 mmol). The reaction was stirred at ambient temperature for 1.5h, water  
10 (25ml) added and the mixture filtered. The solid was washed with water (2x20ml), dried in vacuo and purified by flash chromatography (EtOAc:hexane 1:1) to give the title compound (285mg, 73%) as a powder.

<sup>1</sup>H NMR (mixture of rotamers) δ: 1.2(m, 6H), 2.65(s, 1.5H), 2.85(m, 1.5H), 3.05(m, 1H), 3.2(m, 1H), 4.1(m, 2H), 4.9(m, 2H), 5.55(d, 0.5H), 6.1(d, 0.5H), 7.1(m, 2H), 7.25(m,  
15 3H), 8.5(d, 0.5H), 8.6(d, 0.5H), 12.4(m, 1H); MS *m/z* 506, 508 (M-H).


The following intermediate was made by the process of **Intermediate 20**, using 2,3-dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**) as the amine and (*4S*)-2,2-dimethyl-1,3-dioxolane-4-carboxylic acid potassium salt as the carboxylate source.  
20

**Intermediate 21: 2,3-Dichloro-*N*-[(1*R*,2*R*)-1-[[[(4*S*)-2,2-dimethyl-1,3-dioxolan-4-yl]carbonyl](methyl)amino]-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



5 <sup>1</sup>H NMR (mixture of rotamers): 1.2 (m, 6H), 2.65 (s, 1.5H), 2.85 (s, 1.5H), 3.0 (m, 1H), 3.25 (m, 1H), 4.1 (m, 1H), 4.25 (m, 1H), 4.9 (m, 2H), 5.73 (d, 0.5H), 6.1 (d, 0.5H), 7.1 (m, 2H), 7.25 (m, 3H), 8.62 (d, 0.5H), 8.75 (d, 0.5H), 12.42 (m, 1H); MS *m/z* 506, 508 (M-H).

10 **Intermediate 22 : 2-Chloro-*N*-[(1*R*,2*R*)-1-(methyl[(2*S*)-5-oxotetrahydrofuran-2-yl]carbonyl)amino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**




DIPEA (173  $\mu$ L, 1.0 mmol), (2*S*)-5-oxotetrahydrofuran-2-carboxylic acid (260 mg, 2 mmol) and EDAC (328 mg, 2.0 mmol) were added to a suspension of 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**; 381.5 mg, 1.0 mmol) in anhydrous DMF (5 mL). The reaction was stirred at ambient temperature for 3h. and then diluted with EtOAc (50 mL). The solution was washed with water (4x20 mL), dried (MgSO<sub>4</sub>) and evaporated to give a gum which was triturated with ether to give the title compound (330mg, 72%) as a white solid.

20 <sup>1</sup>H NMR  $\delta$ : 2.2 (m, 2H), 2.5 (, 2H), 2.7 (s, 1.5H), 2.8 (s, 1.5H), 3.0 (m, 1H), 3.3 (m, 1H), 4.85 (m, 1H), 5.5 (m, 1.5H), 6.0 (d, 0.5H), 7.0 (m, 1.5H), 7.1 (m, 0.5H), 7.2 (d, 1H), 7.3 (m, 3H), 8.6 (m, 1.0H), 11.9 (d, 1H); MS *m/z* 456, 458.

The following compound prepared by the process of **Intermediate 22** using 2,3-dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**) as the amine and (2*S*)-5-oxotetrahydrofuran-2-carboxylic acid as the acid.


5

**Intermediate 23: 2,3-Dichloro-*N*-[(1*R*,2*R*)-1-(methyl)[(2*S*)-5-oxotetrahydrofuran-2-yl]carbonyl]amino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



<sup>1</sup>H NMR δ: 2.2 (m, 2H), 2.5 (m, 2H), 2.7 (s, 1.5H), 3.0 (s, 1.0H), 3.3 (m, 1H), 4.9 (m, 1H),  
 10 5.5 (m, 1.5H), 6.05 (d, 0.5H), 7.0 (m, 0.5H), 7.1 (m, 1.5H), 7.3 (m, 3H), 8.7 (m, 1H), 12.44 (d, 1.0H); MS m/z 490, 492.

**Intermediate 24 : 2-Chloro-*N*-((1*R*,2*R*)-1-{methyl[(2*S*)-oxiran-2-ylcarbonyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**




15

EDAC (573 mg, 3 mmol) was added to a stirred suspension of 2-chloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 1**; 381.5mg, 1.0 mmol) and potassium (2*S*)-oxirane-2-carboxylate (378mg, 3mmol) in DMF (5 mL). After stirring for 2h at ambient temperature  
 20 water (20 mL) was added and the resulting solid precipitate was collected by filtration, washed with water and dried under vacuum. MS m/z 416.

The following intermediate was prepared by the method of **Intermediate 24** using 2,3-dichloro-*N*-[(1*R*,2*R*)-1-(methylamino)-2,3-dihydro-1*H*-inden-2-yl]-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide hydrochloride (**Intermediate 16**) as the amine and potassium (2*S*)-oxirane-2-carboxylate as the carboxylic acid.

5


**Intermediate 25: 2,3-Dichloro-*N*-((1*R*,2*R*)-1-{methyl[(2*S*)-oxiran-2-yl]carbonyl}amino)-2,3-dihydro-1*H*-inden-2-yl)-4*H*-thieno[3,2-*b*]pyrrole-5-carboxamide**



MS *m/z* 482.

10

**Intermediate 26: 2-Chloro-*N*-((1*R*,2*R*)-1-[[2-(2,2-dimethyl-1,3-dioxolan-4-yl)carbonyl][2-(tetrahydro-2*H*-pyran-2-yl)oxy]ethyl]amino)-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



- 15 To a solution of 2-chloro-*N*-((1*R*,2*R*)-1-{[2-(tetrahydro-2*H*-pyran-2-yl)oxy]ethyl}amino)-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide (230mg, 0.5 mmol) 2,2-dimethyl-1,3-dioxolane-4-carboxylic acid potassium salt (92mg, 0.5 mmol) in anhydrous DMA (5ml) was added HOBT (68mg, 0.5 mmol) and EDCI (120mg, 0.625mmol). The reaction was stirred at ambient temperature for approximately 24h, water (30ml) added and
- 20 the mixture filtered and dried in vacuo to give the title compound (270mg, 91%) as a powder. MS *m/z* 586, 588 (M-H).


The following intermediates were made by the process of **Intermediate 26**, using 2-Chloro-*N*-((1*R*,2*R*)-1-{[2-(tetrahydro-2*H*-pyran-2-yloxy)ethyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide (**Intermediate 29**) as the amine and (*R*) or (*S*)-2,2-dimethyl-1,3-dioxolane-4-carboxylic acid potassium salt as the carboxylate source.

5

**Intermediate 27: 2-Chloro-*N*-((1*R*,2*R*)-1-{[*(R)*]-2,2-dimethyl-1,3-dioxolan-4-yl)carbonyl}[2-(tetrahydro-2*H*-pyran-2-yloxy)ethyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**


**Intermediate 28: 2-Chloro-*N*-((1*R*,2*R*)-1-{[*(S)*]-2,2-dimethyl-1,3-dioxolan-4-**

10 **yl)carbonyl}[2-(tetrahydro-2*H*-pyran-2-yloxy)ethyl]amino}-2,3-dihydro-1*H*-inden-2-yl)-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide**



Intermediate No	R	MS m/z
27		586, 588
28		586, 588


**Intermediate 29: 2-Chloro-N-((1R,2R)-1-[[2-(tetrahydro-2H-pyran-2-yloxy)ethyl]amino]-2,3-dihydro-1H-inden-2-yl)-6H-thieno[2,3-b]pyrrole-5-carboxamide**



- N*-[(1*R*,2*R*)-1-amino-2,3-dihydro-1*H*-inden-2-yl]-2-chloro-6*H*-thieno[2,3-*b*]pyrrole-5-carboxamide hydrochloride (6.6g, 20 mmol), 2-(2-iodoethoxy)tetrahydro-2*H*-pyran (5.12g, 20mmol) and DIPEA (6.85ml, 40 mmol) in anhydrous DMA (10ml) was heated at 60°C for approximately 24h. The reaction was cooled, EtOAc (150ml) added and washed with water (6 x 50ml), brine (50ml), dried (MgSO<sub>4</sub>) and the volatiles removed under reduced pressure. The residue was purified by flash column chromatography (EtOAc) to give the title compound
- 10 (3.65g, 40%) as a foam.
- <sup>1</sup>H NMR δ: 1.5(m, 6H), 2.15(m, 1H), 2.8(m, 3H), 3.3(m, 3H), 3.65(m, 2H), 4.25(d, 1H), 4.5(m, 2H), 7.0(s, 1H), 7.15(s, 1H), 7.2(m, 3H), 7.3(m, 1H), 8.4(d, 1H), 11.82(m, 1H); MS *m/z* 458, 460 (M-H).

**Claims**

1. A compound of formula (1):



(1)

wherein:

$R^4$  and  $R^5$  together are either  $-S-C(R^6)=C(R^7)-$  or  $-C(R^7)=C(R^6)-S-$ ;

$R^6$  and  $R^7$  are independently selected from hydrogen, halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy, carboxy, carbamoyl,

10 (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy and (1-4C)alkanoyl;

A is phenylene or heteroarylene;

n is 0, 1 or 2;

$R^1$  is independently selected from halo, nitro, cyano, hydroxy, carboxy, carbamoyl,

*N,N*-((1-4C)alkyl)<sub>2</sub>carbamoyl, sulphamoyl, *N*-(1-

15 4C)alkylsulphamoyl, *N,N*-((1-4C)alkyl)<sub>2</sub>sulphamoyl,  $-S(O)_b(1-4C)alkyl$  (wherein b is 0, 1, or

2),  $-OS(O)_2(1-4C)alkyl$ , (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (1-4C)alkanoyl, (1-4C)alkanoyloxy, hydroxy(1-4C)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy and  $-NHSO_2(1-4C)alkyl$ ;

or, when n is 2, the two  $R^1$  groups, together with the carbon atoms of A to which they are

20 attached, may form a 4 to 7 membered saturated ring, optionally containing 1 or 2

heteroatoms independently selected from O, S and N, and optionally being substituted by one or two methyl groups;

one of  $R^2$  and  $R^3$  is selected from  $R_{Na}$ , and the other is selected from  $R_{Nb}$ ;

$R_{Na}$ : (1-3C)alkyl, halo(1-3C)alkyl, dihalo(1-3)alkyl, trifluoromethyl, hydroxy(1-3C)alkyl,

25 dihydroxy(2-3C)alkyl, cyano(1-3C)alkyl (optionally substituted on alkyl with hydroxy),

methoxymethyl, ethoxymethyl, methoxyethyl, methoxymethoxymethyl, dimethoxyethyl,

(hydroxy)(methoxy)ethyl, 5- and 6-membered acetals and mono- and di-methyl derivatives

thereof, (amino)(hydroxy)(2-3C)alkyl, (aminocarbonyl)(hydroxy)(2-3C)alkyl,

(methylaminocarbonyl)(hydroxy)(2-3C)alkyl, (dimethylaminocarbonyl)(hydroxy)(2-3C)alkyl,

(methylcarbonylamino)(hydroxy)(2-3C)alkyl, (methylS(O)<sub>p</sub>)(hydroxy)(2-3C)alkyl (wherein p is 0, 1 or 2);

- R<sub>Nb</sub>: (1-4C)alkyl, halo(1-4C)alkyl, dihalo(1-4C)alkyl, trifluoromethyl, hydroxy(1-4C)alkyl, dihydroxy(2-4C)alkyl, trihydroxy(3-4C)alkyl, cyano(1-4C)alkyl (optionally substituted on alkyl with hydroxy), (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkoxy(1-4C)alkyl, di[(1-4C)alkoxy](2-4C)alkyl, (hydroxy)[(1-4C)alkoxy](2-4C)alkyl, 5- and 6-membered acetals and mono- and di-methyl derivatives thereof, (amino)(hydroxy)(2-4C)alkyl, (aminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, (di(1-4C)alkylaminocarbonyl)(hydroxy)(2-4C)alkyl, ((1-4C)alkylcarbonylamino)(hydroxy)(2-4C)alkyl, ((1-4C)alkylS(O)<sub>p</sub>)(hydroxy)(2-4C)alkyl (wherein p is 0, 1 or 2);
- wherein any alkyl or alkoxy group within any group in R<sub>NA</sub> and R<sub>Nb</sub> may also optionally be substituted on an available carbon atom with a hydroxy group (provided that said carbon atom is not already substituted by a group linked by a heteroatom);
- provided that if R<sup>2</sup> is (1-3C)alkyl or (1-4C)alkyl then R<sup>3</sup> is not (1-4C)alkyl or (1-3C)alkyl; or a pharmaceutically acceptable salt or pro-drug thereof.

2. A compound of formula (1) as claimed in Claim 1, or a pharmaceutically acceptable salt or pro-drug thereof, wherein R<sup>2</sup> is selected from R<sub>NA</sub>, and R<sup>3</sup> is selected from R<sub>Nb</sub>, wherein R<sub>NA</sub> and R<sub>Nb</sub> are as defined in Claim 1.

3. A compound of formula (1) as claimed in Claim 1 or Claim 2, or a pharmaceutically acceptable salt or pro-drug thereof, wherein A is phenylene.

4. A compound of formula (1) as claimed in Claim 1, 2 or 3, or a pharmaceutically acceptable salt or pro-drug thereof, wherein n is 0.


5. A compound of formula (1) as claimed in any one of Claims 1 to 4, or a pharmaceutically acceptable salt or pro-drug thereof, wherein R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen and halo.



6. A compound of formula (1) as claimed in any one of Claims 1 to 4, or a pharmaceutically acceptable salt or pro-drug thereof, wherein  $R^6$  and  $R^7$  are independently selected from hydrogen and chloro.

5 7. A compound of formula (1) as claimed in any one of Claims 1 to 6, or a pharmaceutically acceptable salt or pro-drug thereof, wherein  $R_{Na}$  is selected from (1-4C)alkyl, hydroxy(1-4C)alkyl, and (1-4C)alkoxy(1-4C)alkyl.

8. A compound of formula (1) as claimed in any one of Claims 1 to 7, or a  
10 pharmaceutically acceptable salt or pro-drug thereof, which is a compound of formula (1A):



wherein  $R^1$  to  $R^7$  and  $n$  are as defined in any one of claims 1 to 7.

15 9. A pro-drug of a compound of formula (1) as claimed in any one of Claims 1 to 8, which pro-drug is an in-vivo hydrolysable ester.

10. A pharmaceutical composition which comprises a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1  
20 in association with a pharmaceutically-acceptable diluent or carrier.

11. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, for use in a method of treatment of a warm-blooded animal such as man by therapy.

25

12. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in claim 1, for use as a medicament.

13. A compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in claim 1, for use as a medicament in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.


5

14. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in claim 1, in the manufacture of a medicament for use in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

10


15. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in claim 1, in the manufacture of a medicament for use in the treatment of type 2 diabetes in a warm-blooded animal such as man.

15 16. A process for the preparation of a compound of formula (1) as claimed in claim 1, which process comprises:  
reacting an acid of the formula (2):



(2)

20 or an activated derivative thereof; with an amine of formula (3):



(3)

and thereafter if necessary:

i) converting a compound of the formula (1) into another compound of the formula (1);

25 ii) removing any protecting groups;

iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

# INTERNATIONAL SEARCH REPORT

PCT/GB2004/003546

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/407 C07D495/04 A61P3/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 02/20530 A (FREEMAN SUE ; KENNY PETER (GB); MORLEY ANDREW (GB); WHITTAMORE PAUL (G) 14 March 2002 (2002-03-14) page 36, line 25 - page 39, line 30; examples 149-154	1-16
Y	EP 1 136 071 A (PFIZER PROD INC) 26 September 2001 (2001-09-26) the whole document	1-16
P, X	WO 03/074531 A (WHITTAMORE PAUL ROBERT OWEN ; SIMPSON IAIN (GB); ASTRAZENeca UK LTD (G) 12 September 2003 (2003-09-12) method 10-page 108 page 28, line 16 - page 53, line 30; claims; examples	1-16

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

13 October 2004

Date of mailing of the international search report

20/10/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gavrilu, D

# INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB2004/003546

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0220530	A	14-03-2002	AT 263772 T	15-04-2004
			AU 8283301 A	22-03-2002
			BG 107624 A	30-01-2004
			BR 0113606 A	24-06-2003
			CA 2417594 A1	14-03-2002
			CN 1473163 T	04-02-2004
			CZ 20030616 A3	14-05-2003
			DE 60102710 D1	13-05-2004
			DK 1317459 T3	12-07-2004
			EP 1317459 A1	11-06-2003
			HU 0400784 A2	28-07-2004
			JP 2004508376 T	18-03-2004
			NO 20031024 A	05-03-2003
			NZ 524011 A	27-08-2004
			PT 1317459 T	31-08-2004
			WO 0220530 A1	14-03-2002
			SI 1317459 T1	31-08-2004
			SK 2592003 A3	05-08-2003
			US 2003232875 A1	18-12-2003
EP 1136071	A	26-09-2001	AU 2813001 A	27-09-2001
			CA 2341344 A1	22-09-2001
			EP 1136071 A2	26-09-2001
			HU 0101158 A2	28-02-2002
			JP 2001302546 A	31-10-2001
			NZ 510677 A	25-10-2002
			US 2003004162 A1	02-01-2003
			ZA 200102318 A	20-09-2002
WO 03074531	A	12-09-2003	WO 03074531 A1	12-09-2003